

# A CONSTRUCTIVE VERSION OF TARSKI'S GEOMETRY

MICHAEL BEESON

**Abstract.** Constructivity, in this context, refers to a theory of geometry whose axioms and language are closely related to ruler-and-compass constructions. It may also refer to the use of intuitionistic (or constructive) logic, but the reader who is interested in ruler-and-compass geometry but not in constructive logic, will still find this work of interest. Euclid's reasoning is essentially constructive (in both senses). Tarski's elegant and concise first-order theory of Euclidean geometry, on the other hand, is essentially non-constructive (in both senses), even if we restrict attention (as we do here) to the theory with line-circle continuity in place of first-order Dedekind completeness. Hilbert's axiomatization has a much more elaborate language and many more axioms, but it contains no essential non-constructivities. Here we exhibit three constructive versions of Tarski's theory. One, like Tarski's theory, has existential axioms and no function symbols. We then consider a version in which function symbols are used instead of existential quantifiers. This theory is quantifier-free and proves the continuous dependence on parameters of the terms giving the intersections of lines and circles, and of circles and circles. The third version has a function symbol for the intersection point of two non-parallel, non-coincident lines, instead of only for intersection points produced by Pasch's axiom and the parallel axiom; this choice of function symbols connects directly to ruler-and-compass constructions. All three versions have this in common: the axioms have been modified so that the points they assert to exist are unique and depend continuously on parameters. This modification of Tarski's axioms, with classical logic, has the same theorems as Tarski's theory, but we obtain results connecting it with ruler-and-compass constructions as well. In particular, we show that constructions involving the intersection points of two circles are justified, even though only line-circle continuity is included as an axiom. We obtain metamathematical results based on the Gödel double-negation interpretation, which permit the wholesale importation of proofs of negative theorems from classical to constructive geometry, and of proofs of existential theorems where the object asserted to exist is constructed by a single construction (as opposed to several constructions applying in different cases). In particular, this enables us to import the proofs of correctness of the geometric definitions of addition and multiplication, once these can be given by a uniform construction.

We also show, using cut-elimination, that objects proved to exist can be constructed by ruler and compass. (This was proved in [3] for a version of constructive geometry based on Hilbert's axioms.) Since these theories are interpretable in the theory of Euclidean fields, the independence results about different versions of the parallel postulate given in [5] apply to them; and since addition and multiplication can be defined geometrically, their models are exactly the planes over (constructive) Euclidean fields.

## CONTENTS

1. Introduction	3
2. Hilbert and Tarski	6
2.1. Hilbert	7
2.2. Tarski	8

2.3.	Strict <i>vs.</i> non-strict betweenness and collinearity	9
2.4.	Pasch's axiom	10
2.5.	Sides of a line	11
2.6.	The parallel axiom according to Hilbert and Tarski	12
2.7.	Interpreting Hilbert in Tarski	14
3.	Tarski's theory of straightedge and compass geometry	15
3.1.	Line-circle continuity	16
3.2.	Intersections of circles	18
4.	Tarski's axioms, continuity, and ruler-and-compass constructions	19
4.1.	Segment extension and Euclid I.2	20
4.2.	Degenerate cases of inner Pasch	20
4.3.	Inner Pasch and betweenness	21
5.	Alternate formulations of Tarski's theory	22
5.1.	Continuous Tarski geometry	22
5.2.	Skolemizing Tarski's geometry	23
5.3.	Skolem functions for classical Tarski	23
5.4.	Continuity of the Skolem functions	25
5.5.	Continuity and the triangle circumscription principle	25
6.	A constructive version of Tarski's theory	26
6.1.	Introduction to constructive geometry	26
6.2.	Stability	27
6.3.	Strict and non-strict betweenness	29
6.4.	Intuitionistic Tarski geometry with existential axioms	29
6.5.	Intuitionistic quantifier-free Tarski geometry	30
7.	Perpendiculars, midpoints and circles	31
7.1.	The base of an isosceles triangle has a midpoint	32
7.2.	A lemma of interest only constructively	33
7.3.	Erected perpendiculars from triangle circumscription	33
7.4.	Dropped perpendiculars from line-circle continuity	34
7.5.	Erected perpendiculars from dropped perpendiculars	35
7.6.	Midpoints from perpendiculars	36
7.7.	Gupta's perpendicular construction	39
7.8.	Erected perpendiculars from line-circle continuity and the parallel axiom	40
8.	Other forms of the parallel axiom	41
8.1.	Euclid 5 formulated in Tarski's language	41
8.2.	Tarski's parallel axiom	42
8.3.	Euclid 5 implies Tarski's parallel axiom	43
8.4.	Tarski's parallel axiom implies Euclid 5	44
9.	Uniform perpendicular and uniform rotation	45
9.1.	Uniform perpendicular using line-circle	45
9.2.	Uniform perpendicular without any continuity	46
9.3.	Uniform Reflection	50
9.4.	Equivalence of line-circle and segment-circle continuity	50
9.5.	Representation theorems	51
9.6.	Historical Notes	52
10.	Geometry with terms for the intersections of lines	53

10.1.	Logic of Partial Terms ( <b>LPT</b> )	54
10.2.	A version of Tarski's theory with terms for intersections of lines	55
10.3.	Intersections of lines and the parallel axiom	56
10.4.	Interpreting ruler-and-compass Tarski in intuitionistic Tarski with Skolem functions	60
11.	Relations between classical and constructive geometry	62
11.1.	The double-negation interpretation	63
11.2.	Euclid I.2 revisited	65
11.3.	Hilbert planes and constructive geometry	66
12.	Metatheorems	67
12.1.	Things proved to exist in constructive geometry can be constructed	68
12.2.	Extracting constructions from classical proofs	69
12.3.	Disjunction properties	70
12.4.	Interpretation of Euclidean field theory	71
13.	Circle-circle continuity	72
13.1.	Circle-circle continuity via analytic geometry	72
13.2.	Euclid Book III in Tarski geometry	73
13.3.	The radical axis	75
13.4.	Circle-circle continuity via the radical axis	77
13.5.	Skolem functions for circle-circle continuity	80
14.	Conclusion	81
15.	Listing of axioms for reference	81
15.1.	Classical two-dimensional Tarski geometry	81
15.2.	Intuitionistic Tarski geometry	82
15.3.	Ruler-and-compass Tarski geometry	83

**§1. Introduction.** Euclidean geometry, as presented by Euclid, consists of straightedge-and-compass constructions and rigorous reasoning about the results of those constructions. Tarski's twentieth-century axiomatization of geometry does not bear any direct relation to ruler-and-compass constructions. Here we present modifications of Tarski's theory whose axioms correspond more closely to straightedge-and-compass constructions. These theories can be considered either with intuitionistic (constructive) logic, or with ordinary ("classical") logic. Both versions are of interest.

In [3], we gave an axiomatization of constructive geometry based on a version of Hilbert's axioms (which contain no essential non-constructivities). In [5], we obtained metamathematical results about constructive geometry, and showed that those results do not depend on the details of the axiomatization. In this paper, we focus on formulating constructive geometry in the language and style that Tarski used for his well-known axiomatization of geometry. What is striking about Tarski's theory is its use of only one sort of variables, for points, and

the small number of axioms. Here we give what may be the shortest possible axiomatization of constructive geometry, following Tarski’s example.<sup>1</sup>

In [5], we discussed Euclidean constructive geometry in general terms, and worked informally with a theory that had three sorts of variables for points, lines, and circles. Here, in the spirit of Tarski, we work with a one-sorted theory, with variables for points only. In order to provide terms for points proved to exist, we need some function symbols. Tarski’s axioms have existential quantifiers; we are interested (both classically and constructively) in extensions of the language that provide function symbols to construct points. Three of these symbols are Skolem symbols that correspond immediately to ruler-and-compass constructions: one for extending a segment  $ab$  by another segment  $cd$ , and two for the intersection points of a line and circle. (In our previous work we also had function symbols for the intersection points of two circles; here we will prove that these are not needed, as the intersection points of two circles are already constructible.) Then we need a way to construct certain intersection points of two lines. Such points are proved to exist only by versions of Pasch’s axiom; so one obvious approach is just to provide a Skolem symbol for a suitable version of Pasch’s axiom. (This has been done for decades by people using theorem-provers with Tarski’s axioms.)

However, Tarski’s version of Pasch’s axiom allows “degenerate cases” in which the “triangle” collapses to three points on a line, or the line through the triangle coincides with a side of the triangle. In these cases, the point asserted to exist is not really constructed by intersecting two lines and does not correspond to a ruler-and-compass construction. Therefore, even with classical logic, Tarski’s axioms need some modifications before they really correspond to ruler-and-compass constructions. To start with, we require that the points in Pasch’s axiom be not collinear. Then we have to “put back” the two fundamental axioms about betweenness that Tarski originally had, but which were eliminated when Tarski and his students realized that they followed from the degenerate cases of Pasch. Finally, we have to restrict the segment-extension axiom to extending non-null segments, i.e.  $ab$  with  $a \neq b$ , since extending a null segment is not done by laying a straightedge between two points. More formally, the extension of segment  $ab$  by a non-null segment  $cd$  will not depend continuously on  $a$  as  $a$  approaches  $b$ , while ruler-and-compass constructions should depend continuously on parameters. The resulting modification of Tarski’s classical axioms we call “continuous Tarski geometry”. If we add the function symbols mentioned above, then all those function symbols correspond to ruler-and-compass constructions, and Herbrand’s theorem then tells us that if we can prove  $\forall x \exists y A(x, y)$ , and  $A$  is quantifier-free, then there are finitely many ruler-and-compass constructions  $t_1, \dots, t_n$  such that for each  $x$ , one of the  $t_i(x)$  constructs  $y$  such that  $A(x, y)$ .

We said that ruler-and-compass constructions should depend continuously on parameters, but there is a problem about that: we need to distinguish axiomatically between the two intersection points of a line and a circle. Since lines are given by two distinct points, our solution to this problem is to require that the two intersection points of  $Line(a, b)$  and circle  $C$  occur in the same order on  $L$

---

<sup>1</sup>Readers unfamiliar with Tarski’s geometry may want to begin with [25], which summarizes the axioms of Tarski’s geometry and gives some of their history; but we do give a basic description of Tarski’s axioms in this paper.

as  $a$  and  $b$ . Thus if  $a$  and  $b$  are interchanged, the intersection points given by the two function symbols also are interchanged.

All the changes discussed above make sense and are desirable even with classical logic. They connect the axioms of geometry with ruler-and-compass constructions and, in the case of Pasch's axiom, with its intuitive justification. The degenerate cases of Pasch have nothing to do with triangles and lines; they are really about betweenness relations between points on a single line, so it is philosophically better to formulate the axioms as in continuous Tarski geometry. Having the smallest possible number of axioms is not necessarily the criterion for the best version of a theory.

There is also an issue regarding the best form of the parallel axiom. Historically, several versions have been considered for use with Tarski's theories. Two in particular are of interest: the axiom (A10) that Tarski eventually settled upon, and the "triangle circumscription principle", which says that given three non-collinear points, there is a point  $e$  equidistant from all three (which is then the center of a circle containing the three points). Classically, these two formulations are equivalent, so it is just a matter of personal preference which to take as an axiom. Constructively, the two versions mentioned are also equivalent, as follows from the results of [5] and this paper, but the proof is much lengthier than with classical logic. Euclid's own formulation of the parallel postulate, "Euclid 5", mentions angles, so it requires a reformulation to be expressed in the "points only" language of Tarski's theory; a points-only version of Euclid 5 is given in [5] and repeated below. In [5] it is proved that Euclid 5 is equivalent to the triangle circumscription principle, which is considerably shorter than Euclid 5. We follow Szmielew in adopting the triangle circumscription principle as our parallel axiom, although our results show that we *could* have retained Tarski's version.

There is also "Playfair's axiom", which is the version of the parallel axiom adopted by Hilbert in [13]. That version, unlike all the other versions, makes no existence assertion at all, but only asserts that there cannot exist two different lines parallel to a given line through a given point. This version, making no existence assertion, appears to be constructively weaker than the others, and in [5], it is proved that this is indeed the case.

Our aim in this paper is a constructive version of Tarski's geometry. The changes described above, however, make sense with classical logic and are the primary changes that allow a connection between proofs from Tarski's axioms and ruler-and-compass constructions. If we still use classical logic, proofs in this theory yield a finite number of ruler-and-compass constructions, to be used in the different cases required in the proof. To make the theory constructive, we do just two things more: (1) we use intuitionistic logic instead of classical logic, and (2) we add "stability axioms", allowing us to prove equality or inequality of points by contradiction. It turns out that no more changes are needed. This theory is called "intuitionistic Tarski geometry". As in classical geometry, we can consider it with or without function symbols.

Even though this theory is constructively acceptable, one might not like the fact that the Skolem symbols are total, i.e. everywhere defined; in undefined cases they do not actually correspond to ruler-and-compass constructions. Therefore we also consider a version of Tarski geometry in which the logic is further

modified to use the “logic of partial terms” LPT, permitting the use of undefined terms. In this theory, we replace the Skolem function for Pasch’s axiom by a more natural term  $il(a, b, c, d)$  for the intersection point of  $Line(a, b)$  and  $Line(c, d)$ .

The main difference between constructive and classical geometry is that, in constructive geometry, one is not allowed to make a case distinction. For example, to prove that there always exists a perpendicular to line  $L$  through point  $x$ , we may classically use two different constructions, one of which works when  $x$  is not on  $L$  (a “dropped perpendicular”), and a different construction that works when  $x$  is on  $L$  (an “erected perpendicular”). But constructively, we need a single construction (a “uniform perpendicular”) that handles both cases with one construction. In this paper we show that such uniform constructions can be found, using the Tarski axioms, for perpendiculars, reflections, and rotations. Then the methods of [5] can be used to define addition and multiplication geometrically, as was done classically by Descartes and Hilbert. This shows that every model of the theory is plane over a Euclidean ordered field that can be explicitly constructed.

Having formulated intuitionistic Tarski geometry, we then study its metamathematics, using two logical tools: the Gödel double-negation interpretation, and cut-elimination. The double-negation interpretation is just a formal way of saying that, by pushing double negations inwards, we can convert a classical proof of a basic statement like equality of two points, or incidence of a point on a line, or a betweenness statement, to a constructive proof. (The same is of course not true for statements asserting that something exists.) This provides us with tools for the wholesale importation of certain types of theorems from the long and careful formal development from Tarski’s classical axioms in [19]. But since we modified Tarski’s axioms, to make them correspond better to ruler-and-compass, some care is required in this metatheorem.

Cut-elimination provides us with the theorem that things proved to exist in intuitionistic Tarski geometry can be constructed by ruler and compass. The point here is that they can be constructed by a *uniform* construction, i.e. a single construction that works for all cases. We already mentioned the example of dropped and erected perpendiculars in classical geometry, versus a uniform perpendicular construction in constructive geometry. Using cut-elimination we prove that this feature of constructive proofs, so evident from examples, is a necessary feature of any existence proof in intuitionistic Tarski geometry: an existence proof *always* provides a uniform construction.

On the other hand, our version of Tarski geometry with classical logic, which we call “continuous Tarski geometry”, supports a similar theorem. If it proves  $\forall x \exists y A(x, y)$ , with  $A$  quantifier-free, then there are a finite number (not just one) of ruler-and-compass constructions, given terms of the theory, such that for every  $x$ , one of those constructions produces  $y$  such that  $A(x, y)$ .

Readers familiar with intuitionistic or constructive mathematics will find additional introductory material in § 6.

**§2. Hilbert and Tarski.** It is not our purpose here to review in detail the (long, complicated, and interesting) history of axiomatic geometry, but some

history is helpful in understanding the variety of geometrical axiom systems. We restrict our attention to the two most famous axiomatizations, those of Hilbert and Tarski. Previous work on constructive geometry is discussed in [5].

**2.1. Hilbert.** Hilbert's influential book [13] used the notion of betweenness and the axioms for betweenness studied by Pasch [16]. Hilbert's theory was what would today be called "second-order", in that sets were freely used in the axioms. Segments, for example, were defined as sets of two points, so by definition  $AB = BA$  since the set  $\{A, B\}$  does not depend on the order. Of course, this is a trivial departure from first-order language; but Hilbert's last two axioms, Archimedes's axiom and the continuity axiom, are not expressible in a first-order geometrical theory. On the other hand, lines and planes were regarded not as sets of points, but as (what today would be called) first-order objects, so incidence was an undefined relation, not set-theoretic membership. At the time (1899) the concept of first-order language had not yet been developed, and set theory was still fairly new. Congruence was treated by Hilbert as a binary relation on sets of two points, not as a 4-ary relation on points.

Early geometers thought that the purpose of axioms was to set down the truth about space, so as to ensure accurate and correct reasoning about the one true (or as we now would say, "intended") model of those axioms. Hilbert's book promoted the idea that axioms may have many models; the axioms and deductions from them should make sense if we read "tables, chairs, and beer mugs" instead of "points, lines, and planes." This is evident from the very first sentence of Hilbert's book:

Consider three distinct sets of objects. Let the objects of the first set be called *points* . . . ; let the objects of the second set be called *lines* . . . ; let the objects of the third set be called *planes*.

Hilbert defines segments as pairs of points (the endpoints), although lines are primitive objects. On the other hand, a ray is the set of all points on the ray, and angles are sets consisting of two rays. So an angle is a set of sets of points. Hence technically Hilbert's theory, which is often described as second-order, is at least third-order. Hilbert's language has a congruence relation for segments, and a separate congruence relation for angles. Hilbert's congruence axioms involve the concept of angles: his fourth congruence axiom involves "angle transport" (constructing an angle on a given base equal to a given angle), and his fifth congruence axiom is the SAS triangle congruence principle.

Hilbert's Chapter VII discusses geometric constructions with a limited set of tools, a "segment transporter" and an "angle transporter". These correspond to the betweenness and congruence axioms. Hilbert does not discuss the special cases of line-circle continuity and circle-circle continuity axioms that correspond to ruler-and-compass constructions, despite the mention of "compass" in the section titles of Chapter VII.


Hilbert's geometry contained two axioms that go beyond first-order logic. First, the axiom of Archimedes (which requires the notion of natural number), and second, an axiom of continuity, which essentially says that Dedekind cuts are filled on any line. This axiom requires mentioning a set of points, so Hilbert's

theory with this axiom included is not a “first-order theory” in a language with variables only over points, lines, and circles.

**2.2. Tarski.** Later in the 20th century, when the concept of “first-order theory” was widely understood, Tarski formulated his theory of elementary geometry, in which Hilbert’s axiom of continuity was replaced with an axiom schemata. The set variable in the continuity axiom was replaced by any first-order formula. Tarski proved that this theory (unlike number theory) is complete: every statement in the first-order language can be proved or refuted from Tarski’s axioms. In addition to being a first-order theory, Tarski also made other simplifications. He realized that lines, angles, circles, segments, and rays could all be treated as auxiliary objects, merely enabling the construction of some new points from some given points. Tarski’s axioms are stated using only variables for points. We have listed Tarski’s axioms for reference near the end of this paper, along with the axioms of our constructive version of Tarski geometry, adhering to the numbering of [25], which has become standard.

Tarski replaced Hilbert’s fourth and fifth congruence axioms (angle transport and SAS) with an elegant axiom, known as the five-segment axiom. This axiom is best understood not through its formal statement, but through Fig. 1. The 5-segment axiom says that in Fig. 1, the length of the dashed segment  $cd$  is determined by the lengths of the other four segments in the left-hand triangle. Formally, if the four solid segments in the first triangle are pairwise congruent to the corresponding segments in the second triangle, then the dashed segments are also congruent.

FIGURE 1. Tarski’s 5-segment axiom.  $cd$  is determined.



Tarski’s 5-segment axiom is a thinly-disguised variant of the SAS criterion for triangle congruence. To see this, refer to the figure. The triangles we are to prove congruent are  $dbc$  and  $DBC$ . We are given that  $bc = BC$  and  $db = DB$ . The congruence of angles  $dbc$  and  $DBC$  is expressed in Tarski’s axiom by the congruence of triangles  $abd$  and  $ABD$ , whose sides are pairwise equal. The conclusion, that  $cd = CD$ , give the congruence of triangles  $dbc$  and  $DBC$ . In Chapter 11 of [19], one can find a formal proof of the SAS criterion from the 5-segment axiom. Borsuk-Szmielew also took this as an axiom (see [6], p. 81, Axiom C-5).

An earlier version of Tarski’s theory included as an axiom the “triangle construction theorem”, which says that if we are given triangle  $abc$ , and segment  $AB$  congruent to  $ab$ , and a point  $x$  not on  $Line(A, B)$ , then we can construct

a point  $C$  on the same side of  $Line(A, B)$  as  $x$  such that triangle  $ABC$  is congruent to triangle  $abc$ . It was later realized<sup>2</sup> that this axiom is provable. For example, one can drop a perpendicular from  $c$  to  $Line(a, b)$ , whose foot is the point  $d$  on  $Line(a, b)$ , and then find a corresponding point  $D$  on  $Line(A, B)$ , and then lay off  $dc$  on the perpendicular to  $Line(A, B)$  at  $D$  on the same side of  $Line(A, B)$  as  $x$ , ending in the desired point  $C$ . Of course one must check that this construction can be done and proved correct on the basis of the other axioms. But as it stands, this construction demands a case distinction about the order and possible identity of the points  $d$ ,  $a$ , and  $c$  on  $Line(a, b)$ . Hence, at least this proof of the triangle construction theorem from the axioms of Tarski's theory is non-constructive.

Tarski's early axiom systems also included axioms about betweenness and congruence that were later shown [11] to be superfluous. The final version of this theory appeared in [19]; for the full history see [25].<sup>3</sup> The achievement of Szmielew and Gupta (who are mainly responsible for Part I of [19]) is to develop a really minimal set of axioms for betweenness and congruence.<sup>4</sup> Hilbert's intuitive axioms about betweenness disappeared, leaving only the axiom  $\neg\mathbf{B}(a, b, a)$  and the Pasch axiom and axioms to guarantee that congruence is an equivalence relation.

**2.3. Strict vs. non-strict betweenness and collinearity.** The (strict) betweenness relation is written  $\mathbf{B}(a, b, c)$ . We read this “ $b$  is between  $a$  and  $c$ ”. The intended meaning is that that the three points are collinear and distinct, and  $b$  is the middle one of the three.

Hilbert [13] and Greenberg [10] use strict betweenness, as we do. Tarski [25] used non-strict betweenness. They all used the same letter  $\mathbf{B}$  for the betweenness relation, which is confusing. For clarity we always use  $\mathbf{B}$  for strict betweenness, and introduce  $\mathbf{T}(a, b, c)$  for non-strict betweenness. Since  $\mathbf{T}$  is Tarski's initial, and he used non-strict betweenness, that should be a memory aid. The two notions are interdefinable (even constructively):

DEFINITION 2.1. *Non-strict betweenness is defined by*

$$\mathbf{T}(a, b, c) := \neg(a \neq b \wedge b \neq c \wedge \neg\mathbf{B}(a, b, c))$$

In the other direction,  $\mathbf{B}(a, b, c)$  can be defined as  $\mathbf{T}(a, b, c) \wedge a \neq b \wedge a \neq c$ . The constructive validity of this definition will be discussed at the appropriate time below; here we are still discussing Tarski's classical theory. But we mention this

---

<sup>2</sup>According to [25], Tarski included this principle as an axiom in his first two published axiom sets, but then discovered in 1956-57 with the aid of Eva Kallin and Scott Taylor, that it was derivable; so he did not include it in [24]. (See the footnote, p. 20 of [24].) But Tarski did not publish the proof, and Borsuk-Szmielew take the principle as their Axiom C-7 [6].

<sup>3</sup>Note that the version mentioned in [1] is not the final version used in [19]; inner transitivity for betweenness was eliminated in [19].

<sup>4</sup>We would like to emphasize the important contributions of Gupta, which are important to the development in [19], and are credited appropriately there, but without a careful study one might not realize how central Gupta's results were. These results were apparently never published under Gupta's own name, and still languish in the Berkeley math library in his doctoral dissertation [11]. However, you can get that thesis and others from the Quest Pro database, accessible from most university libraries.

point to emphasize that neither notion is inherently more constructive than the other.

Why then did Tarski choose to use non-strict betweenness, when Hilbert had used strict betweenness? Possibly, as suggested by [25], because this allowed him to both simplify the axioms, and reduce their number. By using  $\mathbf{T}$  instead of  $\mathbf{B}$ , the axioms cover various “degenerate cases”, when diagrams collapse onto lines, etc. Some of these degenerate cases were useful. From the point of view of constructivity, however, this is not desirable. It renders Tarski’s axioms *prima facie* non-constructive (as we will show below). Therefore the inclusion of degenerate cases in the axioms is something that will need to be eliminated in making a constructive version of Tarski’s theories. The same is true even if our only aim is to connect the axioms with ruler-and-compass constructions, while retaining classical logic.

We next want to give a constructive definition of collinearity. Classically we would define this as  $\mathbf{T}(p, a, b) \vee \mathbf{T}(a, p, b) \vee \mathbf{T}(a, b, p)$ . That wouldn’t work as a constructive definition of collinearity, because we have no way to decide in general which alternative might hold, and the constructive meaning of disjunction would require it. In other words, we can know that  $p$  lies on  $Line(a, b)$  without knowing its order relations with  $a$  and  $b$ . But we can find a classically equivalent (yet constructively valid) form by using the law that  $\neg\neg(P \vee Q)$  is equivalent to  $\neg(\neg P \wedge \neg Q)$ . By that method we arrive at

DEFINITION 2.2. *Col(a, b, p) is the formula expressing that a, b, and p lie on a line.*

$$\neg(\neg\mathbf{T}(p, a, b) \wedge \neg\mathbf{T}(a, p, b) \wedge \neg\mathbf{T}(a, b, p))$$

or equivalently, in terms of  $\mathbf{B}$ ,

$$\neg(\neg\mathbf{B}(p, a, b) \wedge \neg\mathbf{B}(a, p, b) \wedge \neg\mathbf{B}(a, b, p) \wedge a \neq p \wedge b \neq p \wedge a \neq b)$$


Note that  $Col(a, b, p)$  only expresses that  $p$  lies on  $Line(a, b)$  if we also specify  $a \neq b$ . We do not put the condition  $a \neq b$  into the definition of  $Col(a, b, p)$  for two reasons: it would destroy the symmetry between the three arguments, and more important, it would cause confusion in comparing our work with the standard reference for Tarski’s theories, namely [19].

**2.4. Pasch’s axiom.** Hilbert’s fourth betweenness axiom is often known as Pasch’s axiom, because it was first studied by Pasch in 1882 [16]. It says that if line  $L$  meets (the interior of) side  $AB$  of triangle  $ABC$  then it meets (the interior of) side  $AC$  or side  $BC$  as well. But Tarski considered instead, two restricted versions of Pasch’s axioms known as “inner Pasch” and “outer Pasch”, illustrated in Fig. 2.

Outer Pasch was an axiom (instead of, not in addition to, inner Pasch) in versions of Tarski’s theories until 1965, when it was proved from inner Pasch in Gupta’s thesis [11], Theorem 3.70, or Satz 9.6 in [19].<sup>5</sup> Outer Pasch appears

<sup>5</sup>But apparently, judging from footnote 4 on p. 191 of [25], Tarski knew as early as 1956-57 that outer Pasch implies inner Pasch; in that footnote Tarski argues against replacing outer Pasch with inner Pasch as an axiom, as Szmielew and Schwabhäuser chose to do. Also on p. 196 of [25], Tarski attributes the idea for the proof of inner Pasch from outer Pasch to specific other people; the history is too long to review here, but he credits only Gupta with the derivation of outer Pasch from inner Pasch.

FIGURE 2. Inner Pasch (left) and outer Pasch (right). Line  $pb$  meets triangle  $acq$  in one side. The open circles show the points asserted to exist on the other side.



as Satz 9.6 in [19]. The proof given in [19] is constructive, and is valid also for strict betweenness. After that, Szmielew chose to take inner Pasch as an axiom instead of outer Pasch, although a footnote in [25] shows that Tarski disagreed with that choice. Gupta’s thesis also contains a proof that outer Pasch implies inner Pasch.

It is not completely clear why Tarski wanted to restrict Pasch’s axiom in the first place, but two good reasons come to mind. First, the restricted forms are valid even in three-dimensional space, so they do not make an implicit dimensional assertion, as the unrestricted Pasch axiom does (it fails in three-space). Second, there is the simpler logical form of inner (or outer) Pasch: unrestricted Pasch needs either a disjunction, or a universal quantifier in the hypothesis, so the condition to be satisfied by the point whose existence is asserted is not quantifier-free and disjunction-free, as it is with inner and outer Pasch. This simplicity of logical form is important for our purposes in constructive geometry, but for Tarski it may just have been a matter of “elegance.”

**2.5. Sides of a line.** The notions of “same side” and “opposite side” of a line will be needed below, and are also of interest in comparing Hilbert’s and Tarski’s geometries. One of Hilbert’s axioms was the *plane separation axiom*, according to which a line separates a plane into (exactly) two regions. Two points  $a$  and  $b$  not on line  $L$  are on opposite sides of  $L$  if  $a \neq b$  and there is a point of  $L$  between  $a$  and  $b$ , i.e., the segment  $ab$  meets  $L$ .

DEFINITION 2.3.


$$\textit{OppositeSide}(a, b, L) := \exists x (on(x, L) \wedge \mathbf{B}(a, x, b))$$

The definition of being on the same side is less straightforward. Hilbert’s definition of  $\textit{SameSide}(a, b, L)$  was that segment  $ab$  does not meet  $L$ . That involves a universal quantifier:

$$\forall x \neg (\mathbf{B}(a, x, b) \wedge on(x, L)).$$

One can get an existential quantifier instead of a universal quantifier by using Tarski's definition, illustrated in Fig. 3:

FIGURE 3. Tarski's definition:  $a$  and  $b$  are on the same side of line  $L$ , as witnessed by point  $c$  on the other side.



DEFINITION 2.4.  $a$  and  $b$  are on the same side of  $L$  if there is some  $c$  such that both  $a$  and  $b$  are on the opposite side of  $L$  from  $c$ . Formally:

$$\text{SameSide}(a, b, L) := \exists c, x, y (\mathbf{B}(a, x, c) \wedge \mathbf{B}(b, y, c) \wedge \text{on}(x, L) \wedge \text{on}(y, L))$$

Another advantage of this definition is that it works in more than two dimensions. It can be proved equivalent to Hilbert's definition above, as is discussed in section 2.7 below.

Hilbert took it as axiomatic that a line divides a plane into two regions. In Tarski's system this becomes a fairly difficult theorem:

THEOREM 2.5 (Plane separation theorem). *If  $p$  and  $q$  are on the same side of line  $L$ , and  $p$  and  $r$  are on opposite sides of  $L$ , then  $q$  and  $r$  are also on opposite sides of  $L$ . Formally,*

$$\text{SameSide}(a, b, L) \wedge \text{OppositeSide}(a, c, L) \rightarrow \text{OppositeSide}(b, c, L)$$

*is provable in neutral constructive geometry (i.e., without using the parallel axiom).*

*Proof.* This is proved in Gupta [11], and also as Satz 9.8 of [19]. The proof follows fairly easily from outer Pasch and the definition of *SameSide*, and occurs in [19] right after the proof of outer Pasch. The proof is completely and unproblematically constructive.

**2.6. The parallel axiom according to Hilbert and Tarski.** As is well-known, there are many propositions equivalent to the parallel postulate in classical geometry. The main point of [5] is to establish which of these versions of the parallel postulate are equivalent in constructive geometry, and which are not. Hilbert's parallel axiom (Axiom IV, p. 25 of [13]) is the version we call Playfair's Axiom, introduced by Playfair in 1729: There cannot be more than one parallel to a given line through a point not on the line. Tarski's axiom A10 as published in [19] is a more complicated statement, classically equivalent. Specifically, it

says that if  $p$  is in the (closed) interior of angle  $\alpha$ , then there exist points  $x$  and  $y$  on the sides of  $\alpha$  such that  $\mathbf{T}(x, p, z)$ . Of course, one cannot mention “interior of angle  $\alpha$ ” directly, so the formulation in Tarski’s language is a bit more complex. Szmielew’s manuscript, on which Part I of [19] is based, took instead the “triangle circumscription principle”, which says that for every three non-collinear points  $a, b, c$ , there exists a point  $d$  equidistant from all three (thus  $d$  is the center of a circle passing through  $a, b$ , and  $c$ , thus circumscribing triangle  $abc$ ).<sup>6</sup>

In [5], we considered the parallel axiom from the constructive point of view, and gave a points-only version of Euclid’s parallel postulate, called “Euclid 5”, as well as a stronger version called the “strong parallel postulate.” These turned out to be constructively equivalent, though the proof requires the prior development of considerable “machinery” based on Euclid 5. We also showed that the triangle circumscription principle is equivalent to the strong parallel postulate, and hence to Euclid 5. In this paper we show that Tarski’s parallel axiom is equivalent to Euclid 5, too. Hence all the versions of the parallel postulate that make an existential assertion turn out to be equivalent.

For the reason of simplicity, we follow Szmielew in using the triangle circumscription principle as the parallel axiom in Tarski’s theories.<sup>7</sup> The center of the circumscribed circle  $abc$  can be constructed with ruler and compass as the intersection point of the perpendicular bisectors of  $ab$  and  $bc$ ; the point of the axiom is that these lines do indeed meet (which without some form of the parallel axiom, they cannot be proved to do). The axiom lends itself well to a points-only theory, since it does not actually mention circles. It merely says there is a point equidistant from the three given points.

Tarski and Givant wrote a letter to Schwabhäuser “around 1978”, which was published in 1998 [25] and has served, in the absence of an English translation of [19], as a common reference for Tarski’s axioms and their history. The letter mentions equivalent versions of the parallel axiom: the two mentioned above and a “Third version of the parallel axiom”, which says that if one connects the midpoints of two sides of a triangle, the connecting segment is congruent to half the third side. In spite of the name “Third version of the parallel axiom”, the letter makes no claim that the different versions are equivalent (in any theory at all). One has to be careful when speaking about “versions of the parallel postulate.” According to [23], p. 51, any statement that holds in Euclidean geometry but not in the standard hyperbolic plane is (classically) equivalent to Euclid’s parallel postulate in Tarski’s geometry with full first-order continuity axioms (Axiom (A11) of [25]). in other words, there are only two complete extensions of neutral geometry with full continuity. But no such thing is true in the theories

---

<sup>6</sup>The triangle circumscription principle is equivalent (with classical logic) to Euclid’s parallel axiom. Euclid IV.5 proves the triangle circumscription principle; the converse implication was first proved by Farkas Bolyai, father of Janos Bolyai, who thought he had proved Euclid’s parallel postulate, but had actually assumed the triangle circumscription principle. See [10], pp. 229–30 and p. 240.

<sup>7</sup>The change in the parallel axiom was apparently one of the “inessential changes” Schwabhäuser introduced in publishing Szmielew’s work. I have not seen Szmielew’s manuscript, but base what I say about it here on [25], page 190.

considered here, which have only line-circle and circle-circle continuity (one as an axiom, and one as a theorem).

Indeed, the “third version” mentioned above is not equivalent to the parallel postulate (in neutral geometry with line-circle and circle-circle continuity), but instead to the weaker assertion that the sum of the angles of every triangle is equal to two right angles. The non-equivalence with the parallel axiom is proved as follows:

**THEOREM 2.6.** *No quantifier-free statement can be equivalent to the parallel axiom in neutral geometry with circle-circle and line-circle continuity.*

*Proof.* We give a model of neutral geometry in which  $M$  (or any quantifier-free formula that is provable with the aid of the parallel axiom) holds, but the parallel axiom fails. Let  $\mathbb{F}$  be a non-Archimedean Euclidean field, and let  $\mathbb{K}$  be the finitely bounded elements of  $\mathbb{F}$ , i.e. elements between  $-n$  and  $n$  for some integer  $n$ . The model is  $\mathbb{K}^2$ . This model is due to Max Dehn, and is described in Example 18.4.3 and Exercise 18.4 of [12], where it is stated that  $\mathbb{K}^2$  is a Hilbert plane, and also satisfies line-circle and circle-circle continuity, since the intersection points with finitely bounded circles have finitely bounded coordinates.

Since  $\mathbb{F}^2$  is a model of geometry including the parallel axiom,  $M$  holds there, and since  $M$  is quantifier free, it holds also in  $\mathbb{K}$ . Yet,  $\mathbb{K}$  is not a Euclidean plane; let  $L$  be the  $x$ -axis and let  $t$  be an infinitesimal. There are many lines through  $(0, 1)$  that are parallel to  $L$  in  $\mathbb{K}$  (all but one of them are restrictions to  $P$  of lines in  $F^2$  that meet the  $x$ -axis at some non finitely bounded point). That completes the proof.

*Discussion.* As remarked above, it follows from Szmielew’s work [23], p. 51, that  $M$  is equivalent to the parallel axiom in Tarki’s geometry with classical logic and the full first-order continuity axiom (A11). The question then arises, how exactly can we use elementary continuity to prove Euclid 5 from  $M$ ? Here is a proof: Assume, for proof by contradiction, the negation of Euclid 5. Then, by elementary continuity, limiting parallels exist (see [10], p. 261). Then Aristotle’s axiom holds, as proved in [12], Prop. 40.8, p. 380. But  $M$  plus Aristotle’s axiom implies Euclid 5 (see [10], p. 220), contradiction, QED.

This proof is interesting because it uses quite a bit of machinery from hyperbolic geometry to prove a result that, on the face of it, has nothing to do with hyperbolic geometry. That is, of course, also true of the proof via Szmielew’s metamathematics. Note that a non-quantifier-free instance of elementary continuity is needed to get the existence of limiting parallels directly; in the presence of Aristotle’s axiom, line-circle continuity suffices (see [10], p. 258), but Aristotle’s axiom does not hold in  $P$ . Finally, the proof of Theorem 2.6 shows that the use of a non-quantifier-free instance of continuity is essential, since quantifier-free instances will hold in Dehn’s model (just like line-circle and circle-circle continuity).

**2.7. Interpreting Hilbert in Tarski.** The fundamental results about betweenness discussed in section 4.3, along with many pages of further work, enabled Szmielew to prove (interpretations of) Hilbert’s axioms in Tarski’s theory. Neither she nor her (posthumous) co-authors pointed this out explicitly in [19],

but it is not difficult to find each of Hilbert's axioms among the theorems of [19] (this has been done explicitly, with computer-checked proofs, in [7]). Here we illustrate by comparing Hilbert's betweenness axioms to Tarski's: Both have symmetry. Hilbert's II,3 can be proved from Tarski's axioms as follows: suppose  $\mathbf{B}(a, b, c)$ . Then  $\neg\mathbf{B}(b, a, c)$ , since if  $\mathbf{B}(b, a, c)$  then  $\mathbf{B}(a, b, a)$ , by inner transitivity and symmetry. Also,  $\neg\mathbf{B}(a, c, b)$ , since if  $\mathbf{B}(a, b, c)$  and  $\mathbf{B}(a, c, b)$ , then  $\mathbf{B}(a, b, a)$  by inner transitivity and symmetry. Hilbert has a "density" axiom (between two distinct points there is a third). This is listed as (A22) in [25], but was never an axiom of Tarski's theory. Density can be proved classically even without line-circle or circle-circle continuity: Gupta ([11], or [19], Satz 8.22) showed that the midpoint of a segment can be constructed without continuity. It is also possible to give a very short direct proof the density lemma:

LEMMA 2.7 (Density). *Suppose  $\mathbf{B}(c, a, p)$ , and  $q$  is a point not on  $ac$ . Then there is a point  $b$  with  $\mathbf{B}(a, b, p)$ .*

*Remark.* Until Ben Richert found this short beautiful proof in 2014, the only proof known to me was to take the third point to be the midpoint.

*Proof.* Extend  $aq$  by  $ca$  to point  $r$ , and extend  $rc$  by  $ca$  to point  $s$ . Apply inner Pasch to  $sprqa$ . The result is point  $b$  with  $\mathbf{B}(a, b, q)$  (and  $\mathbf{B}(q, b, s)$ , but that is irrelevant). That completes the proof.


As discussed above, one can prove in Tarski's system (using the dimension axioms) that Hilbert's and Tarski's definitions of *SameSide* coincide; and Hilbert's plane separation axiom becomes a theorem in Tarski's system.

Hilbert's theory has variables for angles; but in Tarski's theory, angles are given by ordered triples of non-collinear points, and the theory of congruence and ordering of angles has to be developed, somewhat laboriously, but along quite predictable lines, carried out in [19]. Two angles  $abc$  and  $ABC$  are congruent if by adjusting  $a$  and  $c$  on the same rays we can make  $ab = AB$  and  $bc = BC$  and  $ac = AC$ ; or equivalently, if the points on all four rays can be so adjusted; or equivalently, if any adjustment of  $a$  and  $b$  can be matched by an adjustment of  $A$  and  $B$ . The definition of angle ordering is given in [19], Definition 11.27, and the well-definedness depends on an argument by cases using inner and outer Pasch. It would appear on the face of the matter that one needs a more general version of Pasch than inner and outer Pasch. However, that is not actually the case, as we discuss below at the end of Section 10.4. One can therefore use the methods of [19] to construct a conservative extension of Tarski geometry that has variables for angles and directly supports the kind of arguments one finds in Euclid.

It is sometimes possible to reduce theorems about angles directly; in particular it is not necessary to develop the theory of angle ordering to state Euclid's parallel postulate. Here we show how to translate the concept "equal alternating interior angles" into Tarski's language:

**§3. Tarski's theory of straightedge and compass geometry.** Tarski's theory is "elementary" only in the sense that it is first-order. It still goes far

FIGURE 4. Transversal  $pq$  makes alternate interior angles equal with  $L$  and  $K$ , if  $pt = tq$  and  $rt = st$ .



beyond Euclid.<sup>8</sup> To capture Euclid’s geometry, Tarski considered the sub-theory in which the continuity axiom is replaced by “segment-circle continuity”. This axiom asserts the existence of the intersection points of a line segment and a circle, if some point on the segment lies inside the circle and some point on the segment lies outside the circle.

It is this theory that we refer to in the title of this paper as “Tarski’s geometry”.

**3.1. Line-circle continuity.** We now formulate the axiom of line-circle continuity. This tells us when a line and a circle intersect—namely, when there is a point on the line closer (or equally close) to the center than the radius of the circle.<sup>9</sup> But we have not defined inequalities for segments yet, so the formal statement is a bit more complex. Moreover, we have to include the case of a degenerate circle or a line tangent to a circle, without making a case distinction.<sup>10</sup> Therefore we must find a way to express “ $p$  is inside the closed  $\text{Circle}(a, y)$ ”. For that it suffices that there should be some  $x$  non-strictly between  $a$  and  $y$  such that  $ax = ap$ . Since this will appear in the antecedent of the axiom, the “some  $x$ ” will not involve an existential quantifier.

**DEFINITION 3.1.**  $ab < cd$  (or  $cd > ab$ ) means  $\exists x(\mathbf{B}(c, x, d) \wedge ax = ab)$ .  
 $ab \leq cd$  (or  $cd \geq ab$ ) means  $\exists x(\mathbf{T}(c, x, d) \wedge ax = ab)$ , where  $\mathbf{T}$  is non-strict betweenness.

**DEFINITION 3.2.** Let  $C$  be a circle with center  $a$ . Then point  $p$  is **strictly inside**  $C$  means there exists a point  $b$  on  $C$  such that  $ap < ab$ , and  $p$  is **inside**  $C$ , or **non-strictly inside**  $C$ , means  $ap \leq ab$ .

The version of line-circle continuity given in [24] is better described as “segment-circle” continuity:

$$ax = ap \wedge \mathbf{T}(a, x, b) \wedge \mathbf{T}(a, b, y) \wedge ay = aq \rightarrow \exists z(\mathbf{T}(p, z, q) \wedge az = ab)$$


<sup>8</sup>It is confusing that in axiomatic geometry, “elementary” sometimes refers to the elementary constructions, and sometimes to the full first-order theory of Tarski. In this paper we shall not refer again to the full first-order theory.

<sup>9</sup>Note that in spite of the use of the word “circle” the axiom, in the form that only asserts the existence of an intersection point, is valid in  $n$ -dimensional Euclidean space, where it refers to the intersections of lines and spheres.

<sup>10</sup>Avigad *et. al.* count only *transverse* intersection, not tangential intersection, as “intersection.”


This axiom says that if segment  $pq$  meets circle  $C$ , then there is a point  $c$  between  $a$  and  $b$  that lies on the circle. See Fig. 5.

FIGURE 5. Segment-circle continuity.  $p$  is inside the circle,  $q$  is outside, so  $L$  meets the segment  $pq$ .



One may also consider a geometrically simpler formulation of line-circle continuity: if line  $L = \text{Line}(u, v)$  has a point  $p$  inside circle  $C$ , then there is a point that lies on both  $L$  and  $C$ . See Fig. 6.

FIGURE 6. Line-circle continuity.  $p$  is inside the circle, so  $L$  meets the circle.



We consider two versions of this axiom. The weaker version (one-point line-circle) only asserts the existence of one intersection point. The stronger version (two-point line-circle) adds the extra assertion that if  $p \neq b$  (i.e. is strictly inside the circle) then there are two distinct intersection points.

Here are the formal expressions of these axioms

$$\begin{aligned} Col(u, v, p) \wedge u \neq v \wedge \mathbf{T}(a, p, b) &\rightarrow \quad (\text{one-point line-circle}) \\ \exists z (Col(u, v, z) \wedge az = ab) & \end{aligned}$$

$$\begin{aligned} Col(u, v, p) \wedge u \neq v \wedge \mathbf{T}(a, p, b) &\rightarrow \quad (\text{two-point line-circle}) \\ \exists y, z (az = ab \wedge ay = ab \wedge \mathbf{T}(y, p, z) \wedge (p \neq a \rightarrow y \neq z)) & \end{aligned}$$

Classically, we could take a shorter version of two-point line-circle:

$$\begin{aligned} Col(u, v, p) \wedge u \neq v \wedge \mathbf{T}(a, p, b) \wedge p \neq a &\rightarrow \quad (\text{one-point line-circle}) \\ \exists y, z (ay = ab \wedge az = ab \wedge \mathbf{B}(y, p, z)) & \end{aligned}$$

This is classically equivalent to two-point line circle, since the case when  $p = a$  is trivial; but constructively, we cannot make a case distinction whether  $p = a$  or not. The longer form is necessary for a constructive version.

The equivalence of these three continuity axioms, relative to the other axioms of Tarski geometry, is not at all obvious, because

(i) in order to show line-circle implies segment-circle, we need to construct points on the line outside the circle, which requires the triangle inequality. In turn the triangle inequality requires perpendiculars.

(ii) in order to show one-point line-circle implies two-point line-circle, we need to construct the second point somehow. To do that we need to be able to construct a perpendicular to the line through the center. Classically this requires a dropped perpendicular from the center to the line (as the case when the center is on the line is trivial); constructively it requires even more, a “uniform perpendicular” construction that works without a case distinction. But even the former is difficult.

Since two-point line-circle continuity corresponds directly to the uses made (implicitly) of line-circle continuity in Euclid, we adopt it as an axiom of our constructive version(s) of Tarski’s theory. We shall show eventually that all three versions are in fact equivalent, using the other axioms of Tarski’s theory (and not even using any form of the parallel axiom). But this proof rests on the work of Gupta [11], which we will also discuss below.


**3.2. Intersections of circles.** We next give the principle known as circle–circle continuity. should say that if point  $p$  on circle  $K$  lies (non-strictly) inside circle  $C$ , and point  $q$  lies (non-strictly) outside  $C$ , then both intersection points of the circles are defined. This principle would be taken as an axiom, except that it turns out to be derivable from line-circle continuity, so it is not necessary as an axiom. This implication will be proved and discussed fully in § 7, where it will be shown to be true also with intuitionistic logic.<sup>11</sup>

To express that  $p$  is non-strictly inside (or outside) a circle  $C$ , we use the same technique as just above. Namely,  $p$  is inside  $C$  if  $ap = ax$  for some  $x$  non-strictly between  $a$  and a point  $b$  on  $C$ . The situation is illustrated in Fig. 7, where circle  $C$  is given by center  $a$  and point  $b$ , and circle  $K$  is given by center  $c$  and point  $d$ .

---

<sup>11</sup>It is also true that circle-circle continuity implies line-circle continuity. See for example [10], p. 201.

FIGURE 7. Circle-circle continuity.  $p$  is inside  $C$  and  $q$  is outside  $C$ , as witnessed by  $x$  and  $z$ , so the intersection points 1 and 2 exist.



We want this principle to apply even to degenerate circles, and to points that are on  $C$  rather than strictly inside, so we must use  $\mathbf{T}$  rather than  $\mathbf{B}$  to allow  $x = y$  or  $y = z$ , and we must even allow  $a = x = b = z$ .

In order to express this axiom using point variables only, we think of  $K$  as  $\text{Circle}(c, d)$  and  $C$  as  $\text{Circle}(a, b)$ . Then the axiom becomes

$$ap = ax \wedge aq = az \wedge cp = cd \wedge cq = cd \wedge \mathbf{T}(a, x, b) \wedge \mathbf{T}(a, b, z) \rightarrow \\ \exists z_1, z_2 (cz_1 = cd \wedge az_1 = ab \wedge cz_2 = cd \wedge az_2 = ab) \quad (\text{circle-circle})$$

The use of non-strict betweenness  $\mathbf{T}$  allows for the cases when the circles are tangent (either exterior or interior tangency).

<sup>12</sup>

It is not necessary to assert the existence of two distinct intersection points when  $p$  is strictly inside  $C$ , since the second intersection point can be constructed as the reflection of the first in the line connecting the two centers. Then, using the plane separation theorem, one can prove the existence of an intersection point on a given side of the line connecting the centers.

**§4. Tarski's axioms, continuity, and ruler-and-compass constructions.** Two of Tarski's axioms have “degenerate cases”, in the sense that they introduce points that do not depend continuously on the parameters of the axiom. (The two axioms are segment extension, which permits extending a null segment, and inner Pasch, which allows the diagram to collapse to a line.) Even

<sup>12</sup>Proofs of the equivalence of line-circle and circle-circle continuity using Hilbert's axioms (with no continuity and without even the parallel axiom) were found by Strommer [22]. Since these axioms are derivable from (A1)-(A9), as shown by Gupta and Szmielew [19, 7], the equivalence can be proved in (A1)-(A9) (with classical logic). We have not studied the constructivity of Strommer's proof.

using classical logic, we consider this undesirable. We would like to have a formulation of Tarski’s theory that would permit us to use Herbrand’s theorem to show that if  $\exists y A(x, y)$  is provable (where  $x$  stands for several variables, not just one), then there are finitely many ruler-and-compass constructions  $t_1(x), \dots, t_n(x)$  such that for each  $x$ , one of the  $t_i$  constructs the desired  $y$ , i.e.  $A(x, t_i(x))$ . In this section, we discuss how Tarski’s axioms can be slightly modified to eliminate discontinuities.

**4.1. Segment extension and Euclid I.2.** (A4) is the segment construction axiom. Tarski’s version is  $\exists x (\mathbf{T}(q, a, x) \wedge ax = bc)$ . The degenerate case is extending a null segment, i.e. when  $q = a$ ; then the point  $x$  is not uniquely determined, and moreover,  $x$  does not depend continuously on  $q$  as  $q$  approaches  $a$ . One might wonder if  $x = a$ , or in other words  $b = c$  (extending by a null segment) is also a degenerate case, but we do not consider it as degenerate, since there is no discontinuous dependence in that case. Then to avoid degenerate cases, we could consider

$$q \neq a \rightarrow \exists x (\mathbf{T}(q, a, x) \wedge ax = bc) \quad (\text{A4-i})$$

Classically, disallowing  $q = a$  costs nothing, since to extend a null segment  $aa$  by  $bc$ , we just pick any point  $d \neq a$  and extend the non-null segment  $da$  by  $bc$ . Of course, this introduces a discontinuous dependence.

**4.2. Degenerate cases of inner Pasch.** (A7) is inner Pasch; please refer to Fig. 2. This has a degenerate case when  $p = a$  and  $q = b$ , for as  $(p, q)$  approaches  $(a, b)$ , the intersection point  $x$  does not have a unique limit, but could approach any point on  $ab$  or not have a limit at all, depending on how  $(p, q)$  approaches  $(a, b)$ . If  $p = c$  or  $q = c$ , or if  $p = a$  but  $q \neq b$ , or if  $q = b$  but  $p \neq a$ , then there is an obvious choice of  $x$ , so this degenerate case can be removed simply by replacing  $\mathbf{T}$  by  $\mathbf{B}$  in inner Pasch.

Tarski’s version of inner Pasch allows the points  $a, b$ , and  $c$  to be collinear, and this case is technically important, because it allows a number of fundamental theorems about betweenness to be derived that originally were taken as axioms.<sup>13</sup> The point asserted to exist is unique when  $a, b$ , and  $c$  are not collinear; the technical question arises, whether the point can be chosen continuously in the five parameters  $a, b, c, p$ , and  $q$ , in case collinearity is allowed, but the five points are required to be distinct. Some computations (not provided here) show that indeed the point can be continuously chosen.

Nevertheless, we consider the case when  $a, b$ , and  $c$  are collinear to be objectionable, on philosophical grounds. Pasch’s axiom is supposed to justify the construction of certain points by labeling the intersections of lines drawn with a straightedge as actually “existing” points. In the case when the lines coincide, the axiom has no conceptual connection with the idea of intersecting lines, and hence would need some other justification to be accepted as an axiom. If the justification is just that it provides a single axiom from which several intuitively

---

<sup>13</sup>Tarski viewed it as a good thing when the number of axioms could be reduced by using degenerate cases of remaining axioms. We note that in 2013, a further possible reduction in the number of axioms was proved possible by Makarios [15]: interchanging two variables in the conclusion of the five-segment allows the elimination of the symmetry axiom of congruence,  $ab = ba$ .

evident propositions about betweenness can be deduced, that is a distortion of the meaning of the word “axiom.”

Whether or not one gives weight to this philosophical argument, there is a related technical point: we consider below a version of geometry with terms for the intersection points of lines, and we want to be able to use those terms to construct the points shown to exist by Pasch’s axiom. In other words, the problem with Tarski’s too-general version of inner Pasch is that it asserts the existence of points for which there is no ruler-and-compass construction. In that respect, it is unlike any of the other axioms (A1) to (A10), and also unlike the line-circle and circle-circle continuity axioms. This issue reflects in a precise mathematical way the philosophical issue about the collinear case of Pasch’s axiom.

Therefore, we reformulate inner Pasch for continuity, and for constructivity in the sense of ruler-and-compass constructions of the points asserted to exist, as follows:

- We add the hypotheses  $p \neq a$ .
- We add the hypothesis,  $\neg Col(a, b, c)$ .

The resulting axiom is

$$\begin{aligned} & \mathbf{T}(a, p, c) \wedge \mathbf{T}(b, q, c) \wedge p \neq a \wedge \neg Col(a, b, c) \rightarrow \\ & \exists x (\mathbf{B}(p, x, b) \wedge \mathbf{B}(q, x, a)) \end{aligned} \quad (\text{A7-i) strict inner Pasch}$$

Note that we did not require both  $p \neq a$  and  $q \neq b$ . Just one of those is sufficient to allow a ruler-and-compass construction. We do not need two versions, one with  $a \neq p$  and one with  $b \neq q$ , by symmetry. As it turns out, we could use  $\mathbf{B}$  instead of  $\mathbf{T}$  in this axiom and prove the same theorems, as is shown in Section 10.4 below.

**4.3. Inner Pasch and betweenness.** Tarski’s final theory [25] had only one betweenness axiom, known as (A6) or “the identity axiom for betweenness”:

$$\mathbf{T}(a, b, a) \rightarrow a = b.$$

In terms of strict betweenness, that becomes  $\neg \mathbf{B}(a, x, a)$ , or otherwise expressed,  $\mathbf{B}(a, b, c) \rightarrow a \neq c$ . We also refer to this axiom as (A6). The original version of Tarski’s theory had more betweenness axioms (see [25], p. 188). These were all shown eventually to be superfluous in classical Tarski geometry, through the work of Eva Kallin, Scott Taylor, Tarski himself, and especially Tarski’s student H. N. Gupta [11]. These proofs appear in [19]. Here we give the axiom numbers from [25], names by which they are known, and also the theorem numbers of their proofs in [19]:

$$\mathbf{T}(a, b, c) \rightarrow \mathbf{T}(c, b, a) \quad (\text{A14), symmetry, Satz 3.2}$$

$$\mathbf{T}(a, b, d) \wedge \mathbf{T}(b, c, d) \rightarrow \mathbf{T}(a, b, c) \quad (\text{A15), inner transitivity, Satz 3.5a}$$

$$\begin{aligned} & \mathbf{T}(a, b, c) \wedge \mathbf{T}(b, c, d) \wedge b \neq c \rightarrow \\ & \mathbf{T}(a, b, d) \end{aligned} \quad (\text{A16), outer transitivity, Satz 3.7b}$$

$$\begin{aligned} & \mathbf{T}(a, b, d) \wedge \mathbf{T}(a, c, d) \rightarrow \\ & \mathbf{T}(a, b, c) \vee \mathbf{T}(a, c, b) \end{aligned} \quad (\text{A17), inner connectivity, Satz 5.3}$$

$$\begin{aligned} & \mathbf{T}(a, b, c) \wedge \mathbf{T}(a, b, d) \wedge a \neq b \rightarrow \\ & \mathbf{T}(a, c, d) \vee \mathbf{T}(a, d, c) \end{aligned} \quad (\text{A18), outer connectivity, Satz 5.1}$$

The first of these (A14), is a consequence of inner Pasch, formulated with  $\mathbf{T}$ , but the proof uses a degenerate case of inner Pasch, so if we replace inner Pasch by the non-degenerate form (with  $\mathbf{B}$  instead of  $\mathbf{T}$ ), we will (apparently) have to reinstate (A14) as an axiom. The question arises as to whether this is also true of the others. Certainly these cases suffice:

LEMMA 4.1. (A14) and (A15) suffice to prove the collinear case of Tarski's inner Pasch, using (A4-i) and (A7-i) instead of (A4) and (A7). That is,

$$Col(a, b, c) \wedge a \neq b \wedge \mathbf{T}(a, p, c) \wedge \mathbf{T}(b, q, c) \rightarrow \exists x (\mathbf{T}(p, x, b) \wedge \mathbf{T}(q, x, a)).$$

*Proof.* We first note that  $\mathbf{T}(a, b, b)$  follows immediately from the definition of  $\mathbf{T}(a, b, c)$  in terms of  $\mathbf{B}$ .

Since we checked above that the degenerate cases of (A7) are provable, we can assume that all five of the given points are distinct. Since  $Col(a, b, c)$ , we have  $\mathbf{B}(a, b, c) \vee \mathbf{B}(a, c, b) \vee \mathbf{B}(c, a, b)$ .

Case 1,  $\mathbf{B}(a, b, c)$ . Then we take  $x = b$ . We have to prove  $\mathbf{T}(p, b, b) \wedge \mathbf{T}(q, b, a)$ . From  $\mathbf{T}(a, b, c) \wedge \mathbf{T}(b, q, c)$  we have  $\mathbf{T}(a, b, q)$  by (A15). Then  $\mathbf{T}(q, b, a)$  by (A14). Since  $p \neq b$  we have  $\mathbf{T}(p, b, b)$  as shown above. That completes Case 1.

Case 2,  $\mathbf{B}(c, a, b)$ . Then we take  $x = a$ . We have to prove  $\mathbf{T}(p, a, b) \wedge \mathbf{T}(q, a, a)$ . Since  $q \neq a$  we have  $\mathbf{T}(q, a, a)$  as shown above. By symmetry (A14) we have  $\mathbf{T}(a, p, c)$  and  $\mathbf{T}(b, a, c)$ , so by (A15) we have  $\mathbf{T}(b, a, p)$ , so by (A14) again we have  $\mathbf{T}(p, a, b)$  as desired. That completes Case 2.

Case 3,  $\mathbf{B}(a, c, b)$ . Then we take  $x = c$ . We have to prove  $\mathbf{T}(p, c, b) \wedge \mathbf{T}(q, c, a)$ . From  $\mathbf{T}(a, c, b)$  and  $\mathbf{T}(c, q, b)$  we have by (A15)  $\mathbf{T}(a, c, q)$ , whence by (A14),  $\mathbf{T}(q, c, a)$ . From  $\mathbf{T}(a, c, b)$  by (A14), we have  $\mathbf{T}(b, c, a)$ . From  $\mathbf{T}(a, p, c)$  by (A14), we have  $\mathbf{T}(c, p, a)$ . From that and  $\mathbf{T}(b, c, a)$  we have by (A15)  $\mathbf{T}(b, c, p)$ . By (A14) we have  $\mathbf{T}(p, c, b)$  as desired. That completes Case 3, and the proof of the lemma.

**§5. Alternate formulations of Tarski's theory.** In this section we consider some reformulations of Tarski's theories (still using classical logic) that (i) isolate and remove "degenerate cases" of the axioms, and (ii) introduce Skolem functions to achieve a quantifier-free axiomatization, and (iii) introduce additional axioms to make the intersection points of lines and circles, or circles and circles, depend continuously on the (points determining the) lines and circles.

**5.1. Continuous Tarski geometry.** Let "continuous Tarski geometry" refer to classical Tarski geometry with two-point line-circle continuity, with the following modifications:

- (A4-i) instead of (A4) (extending non-null segments)
- (A7-i) (strict inner Pasch) instead of (A7). That is, use  $\mathbf{B}$  instead of  $\mathbf{T}$  in Pasch, and require  $\neg Col(a, b, c)$  and  $p \neq a$ .
- Take (A14) and (A15) as axioms (symmetry and transitivity of betweenness)
- Use the triangle circumscription principle (A10<sub>3</sub>) for the parallel axiom

The reason for the name "continuous Tarski geometry" will be apparent eventually, when we show what seems intuitively obvious: that Skolem functions for these axioms can be implemented by ruler-and-compass constructions.

**THEOREM 5.1.** *Continuous Tarski geometry has the same theorems as Tarski geometry.*

*Proof.* To extend a null segment  $bb$  by  $cd$ , first select any point  $a$  different from  $b$ , then extend  $ab$  by  $cd$ . Hence the restriction to (A4-i) costs nothing. By Lemma 4.1, the restriction to the non-collinear and non-degenerate case of (A7) is made up for by the inclusion of (A14) and (A15) as axioms. That completes the proof of the theorem.

**5.2. Skolemizing Tarski's geometry.** Since Tarski's axioms are already in existential form, one can add Skolem functions to make them quantifier-free. Perhaps the reason why Tarski did not do so, is his desire that there should be just one model of his theory over the real plane  $\mathbb{R}^2$ . If one introduces Skolem functions for the intersection points of two circles, then those Skolem functions can be interpreted quite arbitrarily, unless one also adds further axioms to guarantee their continuity, and even then, one has a problem because those Skolem functions will be meaningless (have arbitrary values) when the circles do not intersect. Tarski did not have line-circle continuity, but the same problem arises with Skolem functions for inner Pasch, when the hypotheses are not satisfied.

The problem can be seen in a simpler context, when we try to axiomatize field theory with a function symbol  $i(x)$ , the official version of  $x^{-1}$ . The point is that 0 has no multiplicative inverse, yet Skolem functions are total, so  $i(0)$  has to denote *something*. We phrase the axiom as  $x \neq 0 \rightarrow x \cdot i(x) = 1$ , so we can't prove  $0 \cdot i(0) = 1$ , which is good, since we *can* prove  $0 \cdot i(0) = 0$ . In spite of this difficulty, the theory with Skolem functions is a conservative extension of the theory without Skolem functions, as one sees (for theories with classical logic) from the fact that every model of the theory without can be expanded by suitably interpreting the Skolem function symbols. We return below to the question of how this works for intuitionistic theories in Lemma 6.4 below.

Papers on axiomatic geometry often use the phrase "constructive theory" to mean one with enough function symbols to be formulated with quantifier-free axioms. While this is not sufficient to imply that a theory is "constructive" in the sense of being in accordance with Bishop's constructive mathematics (or another branch of constructive mathematics), it is a desirable feature, in the sense that a constructive theory should provide terms to describe the objects it can prove to exist. In finding a constructive version of Tarski's theories, therefore, we will wish to produce a version with function symbols corresponding to ruler and compass constructions. In order to compare the constructive theory with Tarski's classical theory, we will first consider a Skolemized version of Tarski's theory, with classical logic.

**5.3. Skolem functions for classical Tarski.** One introduces Skolem functions and reformulates the axioms to be quantifier-free. But we want these Skolem functions to be meaningful as ruler-and-compass constructions. Hence, we do not Skolemize Tarski's theory as he gave it, but rather the modified version we called "continuous Tarski geometry." The axioms are listed for reference at the end of the paper; here we just give a list of the Skolem functions:

- $ext(a, b, c, d)$  is a point  $x$  such that for  $a \neq b$ , we have  $\mathbf{T}(a, b, x) \wedge bx = cd$ .

- $ip(a, p, c, b, q)$  is the point asserted to exist by inner Pasch (see Fig 2), provided  $a$ ,  $b$ , and  $c$  are not collinear, and no two of the five points are equal.
- Three constants  $\alpha$ ,  $\beta$ , and  $\gamma$  for three non-collinear points. (In this paper we consider only plane geometry, for simplicity.)
- $center(a, b, c)$  is a point equidistant from  $a$ ,  $b$ , and  $c$ , provided  $a$ ,  $b$ , and  $c$  are not collinear.
- $ilc_1(a, b, c, d)$  and  $ilc_2(a, b, c, d)$  for the two intersection points of  $Line(a, b)$  and  $Circle(c, d)$ , the circle with center  $c$  passing through  $d$ .

The function  $center$  is needed to remove the existential quantifier in Szmielew’s parallel axiom (A10<sub>2</sub>), which says that if  $a$ ,  $b$ , and  $c$  are not collinear, there exists a circle through  $a$ ,  $b$ , and  $c$ . For the version (A10) of the parallel axiom used in [19], we would need two different Skolem functions. The points asserted to exist by that version are not unique and do not correspond to any natural ruler-and-compass construction, which is another reason to prefer triangle circumscription as the parallel axiom.

The question arises, what do we do about “undefined terms”, e.g.,  $ilc_1(a, b, c, d)$  when the line and circle in question do not actually meet? One approach is to modify the logic, using the “logic of partial terms”, introducing a new atomic statement  $t \downarrow$  (read “ $t$  is defined”) for each term  $t$ . In Tarski’s geometry as described here, that is not necessary, since we can explicitly give the conditions for each term to be defined. In that way,  $t \downarrow$  can be regarded as an abbreviation at the meta-level, rather than an official formula. We write the formula as  $(t \downarrow)^\circ$  to avoid confusion and for consistency of notation with another section below.

**DEFINITION 5.2.** *When the arguments to the Skolem functions are variables or constants, we have*

$$\begin{aligned} (ext(a, b, c, d) \downarrow)^\circ &:= a \neq b \\ ip(a, p, c, b, q) \downarrow)^\circ &:= \mathbf{T}(a, p, c) \wedge \mathbf{T}(b, q, c) \wedge a \neq p \wedge \neg Col(a, b, c) \\ (center(a, b, c) \downarrow)^\circ &:= \neg Col(a, b, c) \end{aligned}$$

*If the arguments  $a$ ,  $b$ ,  $c$ ,  $d$  are not variables or constants, then we need to add (recursively) the formulas expressing their definedness on the right.*

In addition to the obvious “Skolem axioms” involving these function symbols, we will need additional axioms to ensure that the two intersection points of a line and circle are distinguished from each other (except when the intersection is of a circle and a tangent line), and that the intersection points depend continuously on the (points determining the) lines and circles.

We discuss the two points of intersection of  $Line(a, b)$  and  $Circle(c, d)$ , which are denoted by  $ilc_1(a, b, c, d)$  and  $ilc_2(a, b, c, d)$ . We want an axiom asserting that these two points occur on  $Line(a, b)$  in the same order as  $a$  and  $b$  do; that axiom serves to distinguish the two points and ensure that they depend continuously on  $a, b, c$ , and  $d$ . To that end we need to define  $SameOrder(a, b, c, d)$ , assuming

$a \neq b$  but allowing  $c = d$ . This can be done as follows:

$$\begin{aligned} \text{SameOrder}(a, b, c, d) &:= (\mathbf{T}(c, a, b) \rightarrow \neg\mathbf{B}(d, c, a)) \\ &\quad \wedge (\mathbf{T}(a, c, b) \rightarrow \neg\mathbf{B}(d, c, b)) \\ &\quad \wedge (\mathbf{T}(a, b, c) \rightarrow \mathbf{T}(a, c, d)) \end{aligned}$$

The axiom in question is then

$$\text{SameOrder}(a, b, \text{ilc}_1(a, b, c, d), \text{ilc}_2(a, b, c, d)).$$

**5.4. Continuity of the Skolem functions.** We will investigate what additional axioms are necessary to guarantee that the Skolem functions are uniquely defined and continuous. Unless we are using the logic of partial terms, technically Skolem functions are total, in which case we can't avoid some arbitrariness in their values, but when their "definedness conditions" given above are satisfied, we expect them to be uniquely defined and continuous. This will be important for metatheorems about the continuous dependence on parameters of things proved constructively to exist; but we think it is also of interest even to the classical geometer.

Evidently for this purpose we should use the version of the axioms that has been sanitized of degenerate cases. Thus, *ext* only Skolemizes axiom (A4-i), for extending non-degenerate segments, and *ip* only Skolemizes axiom (A7-i) rather than A7. These Skolem functions will then be uniquely defined (and provably so).

LEMMA 5.3. *The terms  $\text{ilc}_1(a, b, c, d)$  and  $\text{ilc}_2(a, b, c, d)$  are provably continuous in  $a, b, c,$  and  $d$  (when their definedness conditions hold).*

*Proof.* Once we have defined multiplication and addition, this proof can be carried out within geometry, using ordinary algebraic calculations. It is very much easier to believe that these (omitted) proofs can be carried out, than it is to actually get a theorem-prover or proof-checker to do so. See [4] for a full discussion of the issues involved.

THEOREM 5.4 (Continuity of inner Pasch). *Tarski's geometry, using axioms (A4-i) and (A7-i), proves the continuity of  $\text{ip}(a, p, c, b, q)$  as a function of its five parameters, when the hypotheses of inner Pasch are satisfied.*

*Remark.* If we use axiom (A7), without the modifications in (A7-i), then  $\text{ip}(a, p, c, b, q)$  is not continuous as  $(p, q)$  approaches  $(a, b)$ , as discussed above.

*Proof.* This also can be carried out by introducing coordinates and making ordinary algebraic computations within Tarski geometry.

**5.5. Continuity and the triangle circumscription principle.** Above we have given the triangle circumscription principle with the hypothesis that  $a, b,$  and  $c$  are non-collinear (and hence distinct) points. What happens when that requirement is relaxed? If  $a$  and  $b$  are allowed to approach each other without restriction on the direction of approach, then  $\text{center}(a, b, c)$  does not depend continuously on its parameters. But if  $a$  and  $b$  are restricted to lie on a fixed line  $L$  (as is the case when using triangle circumscription to define multiplication as Hilbert did), then as  $a$  approaches  $b$  (both remaining away from  $c$ ), the circle

through  $a$ ,  $b$ , and  $c$  nicely approaches the circle through  $a$  and  $c$  that is tangent to  $L$  at  $a$ . The *strong triangle circumscription principle* says that there is a term  $C(a, b, c, p, q)$  such that when  $a$  and  $b$  lie on  $L = \text{Line}(p, q)$  and  $c$  does not lie on  $L$ , then  $e = C(a, b, p, q)$  is equidistant from  $a$ ,  $b$ , and  $c$ , and moreover, if  $a = b$  then  $ea$  is perpendicular to  $L$  at  $a$  (i.e., the circle is tangent to  $L$  at  $a$ ). In [5], it is shown how to construct the term  $C$ , using segment extensions and the uniform perpendicular; so this construction can be carried out in Tarski geometry with Skolem functions.

**§6. A constructive version of Tarski’s theory.** Finally we are ready to move from classical to intuitionistic logic. Our plan is to give two intuitionistic versions of Tarski’s theory, one with function symbols as in the Skolemized version above, and one with existential axioms as in Tarski’s original theory. The underlying logic will be intuitionistic predicate logic. We first give the specifically intuitionistic parts of our theory, which are very few in number. We do not adopt decidable equality ( $a = b \vee a \neq b$ ), nor even the substitute concept of “apartness” introduced by Brouwer and Heyting (and discussed below), primarily because we aim to develop a system in which definable terms (constructions) denote continuous functions, but also because we wish to keep our system closely related to Euclid’s geometry, which contains nothing like apartness.

**6.1. Introduction to constructive geometry.** Here we discuss some issues particular to geometry with intuitionistic logic. The main point is that we must avoid case distinctions in existence proofs. What one has to avoid in constructive geometry is not proofs of equality or inequality by contradiction, but rather constructions (existence proofs) that make a case distinction. For example, classically we have two different constructions of a perpendicular through point  $p$  to line  $L$ , one for when  $p$  is not on  $L$ , and another for when  $p$  is on  $L$ . Pushing a double negation through an implication, we only get not-not a perpendicular exists, which is not enough. To constructivize the theorem, we have to give a uniform construction of the perpendicular, which works without a case distinction. (Such a construction is given in [5].)

In particular, in order to show that the models of geometry are planes over Euclidean fields, we need to define addition and multiplication by just such uniform constructions, without case distinctions about the sign of the arguments. The classical definitions due to Descartes and Hilbert do depend on such case distinctions; in [5] we have given uniform definitions; here we check that their properties can be proved in intuitionistic Tarski geometry. To actually carry out the complete development directly would be a project of about the length and scope of Szmielew’s comparable development of classical geometry from Tarski’s axioms, in Part I of [19]. Therefore it is important that the double-negation interpretation can be made to carry the load.

We mention here two principles which are not accepted by all constructivists. Here  $x < y$  refers to points on a fixed line  $L$ , and can be defined in terms of betweenness.

$$\begin{array}{ll} \neg\neg x > 0 \rightarrow x > 0 & \text{(Markov’s principle)} \\ x \neq 0 \rightarrow x < 0 \vee x > 0 & \text{(two-sides)} \end{array}$$

Markov's principle follows from the stability of betweenness and is a fundamental principle of constructive geometry. It allows us to avoid distinguishing more than one sense of inequality between points. Geometry without it would be much more complicated. The principle "two-sides" is closely related to "a point not on a given line is on one side or the other of the line". (Here the "line" could be the  $y$ -axis, i.e. a line perpendicular to  $L$  at the point 0.) This principle is not needed in the formalization of Euclid, or the development of the geometrical theory of arithmetic, and as we will show, it is not a theorem of intuitionistic Tarski geometry.

One might consider adopting two-sides as an axiom, on grounds similar to those sometimes used to justify Markov's principle or apartness, namely that if we "compute  $x$  to sufficient accuracy we will see what sign it has." That justification applies only to the model of computable reals, not to various more general intuitionistic models of sequences generated by free choices of approximations to points. Brouwer argued against this principle in one of his later papers [8] on those grounds; and our development of constructive geometry shows that it is not needed for the usual theorems, including the geometric definitions of addition and multiplication. In our opinion, not only is it unnecessary, it is also constructively undesirable, as the choice of which disjunct holds cannot depend continuously on  $x$ , so anyone claiming its validity must make some assumptions about how points are "given", e.g. by a computable sequence of rational approximations; we do not want to make such assumptions.

On the other hand, the following principle *has* been accepted by all constructivists in the past who considered geometry:

$$a < b \rightarrow x < b \vee a < x \quad (\text{apartness})$$

It turns out that apartness is completely unnecessary for the formalization of Euclid, and is not a theorem of intuitionistic Tarski geometry. The desire to use apartness probably arose from an unwillingness to the trichotomy law of order, and to find some replacement for it. In our work, the law of trichotomy of order is replaced by the stability of equality and betweenness. If we want to formalize one of Euclid's proofs where two points are proved equal by contradiction (consider III.4 for a specific example), the proof in Euclid shows  $\neg a \neq b$ ; in other words  $\neg\neg a = b$ . So, by the stability of equality,  $a = b$ . The trichotomy law can similarly be double negated, each case but one shown contradictory, and the final double negation removed by stability. That is the fundamental reason why apartness is not needed in constructive geometry.

**6.2. Stability.** The word "stable" is applied to a predicate  $Q$  if  $\neg\neg Q \rightarrow Q$ . Our intuitionistic versions of Tarski geometry will all have axioms of stability for the basic predicates. That is, we will include the axioms

$$\begin{aligned} \neg a \neq b &\rightarrow a = b \\ \neg\neg\mathbf{B}(a, b, c) &\rightarrow \mathbf{B}(a, b, c) \\ \neg\neg ab = cd &\rightarrow ab = cd \end{aligned}$$

In this section we justify accepting these axioms. Our intuition is that there is nothing asserting existence in the meaning of equality, congruence, or betweenness; hence assertions of equality, congruence, or betweenness can be constructively proved by contradiction. There are many examples in Euclid<sup>14</sup> where Euclid argues that two points, differently constructed, must coincide; such examples use the stability of equality. Similarly, if point  $x$  lies on  $Line(a, b)$ , we may wish to argue by cases as to its position on the line relative to  $a$  and  $b$ . We double-negate the disjunction of the five possible positions, argue each case independently, and arrive at the double negation of the desired conclusion. As long as what we are proving is a betweenness, congruence, or equality, stability allows us to remove the double negation and reach the desired conclusion.

We explain this point with more detail, for those inexperienced with intuitionistic reasoning: Suppose  $P \rightarrow Q$ , and  $R \rightarrow Q$ . Then classically,  $(P \vee R) \rightarrow Q$ , which does not follow intuitionistically. But intuitionistically, we still have  $\neg\neg(P \vee R) \rightarrow \neg\neg Q$ , since if  $\neg Q$  then  $\neg P$  and  $\neg R$ , which is  $\neg(P \vee R)$ , contradicting  $\neg\neg(P \vee R)$ . Now if  $Q$  is stable we can still conclude  $Q$ .

What we are not allowed to do, constructively, is argue by cases for an existential conclusion, using a different construction for each case. (In the previous paragraph, if  $Q$  begins with  $\exists$ , then  $Q$  will not be stable.) This observation makes it apparent why the constructivization of geometry hinges on the successful discovery of *uniform* constructions, continuous in parameters.

As we mentioned above, angles can be defined in Tarski's theory, and one can show that the equality and ordering of angles is stable. That is,

$$\neg\neg\alpha < \beta \rightarrow \alpha < \beta$$

for angles  $\alpha$  and  $\beta$ . Thus, when Euclid wants to prove  $\alpha = \beta$ , and says, if not, then one of them is greater; let  $\alpha > \beta$ , and so on, the reasoning is constructive, because we have

$$\neg\neg(\alpha < \beta \vee \alpha = \beta \vee \beta < \alpha)$$

and if  $\alpha < \beta$  and  $\beta < \alpha$  lead to contradictions, then  $\neg\neg\alpha = \beta$ , whence by stability,  $\alpha = \beta$ . Similarly if what is to be proved is an inequality of angles.

Julien Narboux pointed out that the stability of equality can be derived from the stability of congruence:

LEMMA 6.1. *With the aid of axioms A1 and A3, stability of congruence implies stability of equality.*

*Proof.* Suppose  $\neg a = b$ . We want to prove  $a = b$ . By A3, it suffices to prove  $ab = aa$ . By the stability of congruence, we may prove this by contradiction. Suppose, for proof by contradiction, that  $ab \neq aa$ . We claim  $a \neq b$ . To prove it, suppose  $a = b$ . Then from  $ab \neq aa$  we obtain  $ab \neq ab$ , contradicting A1. Therefore  $a \neq b$ . But that contradicts the hypothesis  $\neg a = b$  from the first line of the proof. That completes the proof of the lemma.

We could therefore drop stability of equality as an axiom, but we retain it anyway, because of its fundamental character, and to emphasize that it is perhaps even more fundamental than the facts expressed in A1 and A3.

<sup>14</sup>Just to mention one, Euclid III.4

*Stability of incidence.* Tarski's theory has variables for points only, so when we discuss lines, implicitly each line  $L$  is given by two points,  $L = \text{Line}(a, b)$ . When we say point  $x$  lies on line  $L$ , that abbreviates

$$\neg\neg(\mathbf{T}(x, a, b) \vee \mathbf{T}(a, x, b) \vee \mathbf{T}(a, b, x)).$$

Since logically,  $\neg\neg\neg P$  is equivalent to  $\neg P$ , the relation  $x$  lies on  $L$  is stable. (Four negations is the same as two negations.) We refer to this as the “stability of incidence.” When the definition of incidence is expanded, this is seen to be a logical triviality, not even worth of the name “lemma.” But when working in Tarski's theories with less than complete formality, we do mention lines and incidence and justify some proof steps by the “stability of incidence.” In other words, we are allowed to prove that a point  $x$  lies on a line  $L$  by contradiction.

**6.3. Strict and non-strict betweenness.** Should we use strict or non-strict betweenness in constructive geometry? The answer is, it doesn't matter much, because of the stability of  $\mathbf{B}$ . What we do officially is use strict betweenness  $\mathbf{B}$ , and regard  $\mathbf{T}$  as defined by

$$\mathbf{T}(a, b, c) := \neg(a \neq b \wedge b \neq c \wedge \neg\mathbf{B}(a, b, c)).$$

We could also have taken  $\mathbf{T}$  as primitive and defined  $\mathbf{B}$  by

$$\mathbf{B}(a, b, c) := \mathbf{T}(a, b, c) \wedge a \neq b \wedge b \neq c.$$

**6.4. Intuitionistic Tarski geometry with existential axioms.** The language of this theory takes strict betweenness  $\mathbf{B}(a, b, c)$  as primitive, and  $\mathbf{T}(a, b, c)$  will then be a defined concept, given by Definition 2.1. Some of the axioms will be “unmodified” from Tarski's theory, by which we mean that the only change is to define  $\mathbf{T}$  in terms of  $\mathbf{B}$ . The other modifications are as follows:

- Modify Axiom A4 (segment extension) so only non degenerate segments are extendable:  $q \neq a \rightarrow \exists x (\mathbf{T}(q, a, x) \wedge ax = bc)$  (A4-i)

- Axiom (A6) become  $\neg\mathbf{B}(a, b, a)$ .
- Replace inner Pasch (A7) by (A7-i), which requires  $\neg\text{Col}(a, b, c)$  and  $a \neq p$ :

$$\mathbf{T}(a, p, c) \wedge \mathbf{T}(b, q, c) \wedge a \neq p \wedge \neg\text{Col}(a, b, c) \rightarrow \exists x (\mathbf{B}(p, x, b) \wedge \mathbf{B}(q, x, a))$$

- Add (A14-i), the symmetry axiom for betweenness:

$$\mathbf{B}(a, b, c) \rightarrow \mathbf{B}(c, b, a)$$

- Add (A15-i), inner transitivity of betweenness:

$$\mathbf{B}(a, b, d) \wedge \mathbf{B}(b, c, d) \rightarrow \mathbf{B}(a, b, c)$$

- Use Szmielew's preferred version of the parallel postulate, the triangle circumscription principle,

$$\forall a, b, c (\neg\text{Col}(a, b, c) \rightarrow \exists d (ad = bd \wedge ad = cd)).$$

- Add two-point line-circle continuity.
- Use intuitionistic logic only.

- add the stability of equality, betweenness, and congruence:

$$\begin{aligned} \neg a \neq b &\rightarrow a = b \\ \neg \mathbf{B}(a, b, c) &\rightarrow \mathbf{B}(a, b, c) \\ \neg \neg ab = cd &\rightarrow ab = cd \end{aligned}$$

The resulting theory is called “intuitionistic Tarski geometry”, or “intuitionistic Tarski geometry with existential axioms.” Another way of describing it is: restrict continuous Tarski geometry to intuitionistic logic, and add the stability axioms for equality, betweenness, and equidistance, use the triangle circumscription principle for the parallel axiom, and add two-point line-circle continuity. We use the phrase “intuitionistic Tarski geometry without any continuity” to refer to the theory obtained by dropping the line-circle continuity axiom.

**THEOREM 6.2.** *Intuitionistic Tarski geometry plus classical logic is equivalent to Tarski geometry (with or without line-circle continuity).*

*Proof.* This follows from Theorem 5.1, Since intuitionistic Tarski geometry is continuous Tarski geometry with intuitionistic logic and stability axioms, it is classically equivalent to continuous Tarski geometry. But by Theorem 5.1, that theory is classically equivalent to Tarski geometry. That completes the proof.

**6.5. Intuitionistic quantifier-free Tarski geometry.** We can use the same Skolem functions as for the classical theory, since we already made the necessary restrictions to the Skolem functions for segment extension and Pasch’s axioms. For the same reason, the conditions for definedness of Skolem terms are not changed.

**LEMMA 6.3.** *For every term  $t$  of intuitionistic quantifier-free Tarski geometry, the sentence  $\neg \neg t \downarrow \rightarrow t \downarrow$  is provable.*

*Proof.* By induction on the complexity of  $t$ , using the stability of  $\mathbf{B}$ ,  $E$ , and equality for the base case.

Since the conditions for the definedness of Skolem terms are definable, there is no logical problem about using (total) Skolem functions in this intuitionistic theory, without modifying the logic, which is the ordinary intuitionistic first-order predicate calculus. However, there might be a philosophical problem, as one might ask, what is the intended interpretation of those total Skolem symbols? One cannot specify a total (everywhere defined) construction to interpret, for example, the Skolem symbol for inner Pasch. Therefore it is more philosophically correct to use the “logic of partial terms”, which is explained in another section below. However, it is possible to consider the Skolem symbols as mere syntactic tools, which, even if not meaningful, at least cause no unwanted deductions, according to the following lemma:

**LEMMA 6.4.** *[Conservativity of Skolem functions] Suppose intuitionistic Tarski with Skolem functions proves a theorem  $\phi$  that does not contain Skolem functions. Then intuitionistic Tarski (with existential quantifiers and no Skolem functions) also proves  $\phi$ . In fact, the same is true of any intuitionistic theory whose axioms before Skolemization have the form  $P(x) \rightarrow \exists y Q(x, y)$ , with  $P$  quantifier-free.*

*Proof.* Consider a Skolem symbol with axiom  $P(x) \rightarrow Q(x, f(x))$ , Skolemizing the axiom  $P(x) \rightarrow \exists y Q(x, y)$ . The corresponding lemma for classical theories needs no restriction on the form of  $P$ ; one simply shows that every model of the theory without Skolem functions can be expanded to a model of the theory with Skolem functions. The interpretation of the values of a Skolem symbol, say  $f(b)$  are just taken arbitrarily when  $P(b)$  is not satisfied. Then one appeals to the completeness theorem. One can use the Kripke completeness theorem to make a similar argument for theories with intuitionistic logic; but in general one cannot define  $f(b)$  at a node  $M$  of a Kripke model where  $P(b)$  fails, because  $P(b)$  might hold later on, and worse, there might be nodes  $M_1$  and  $M_2$  above  $M$  at which different values of  $y$  are required, so there might be no way to define  $f(b)$  at  $M$ . That cannot happen, however, if  $P$  is quantifier-free, since then, if  $P(b)$  does not hold at  $M$ , it also doesn't hold at any node above  $M$ . Hence if  $P$  is quantifier free, we can complete the proof, using Kripke completeness instead of Gödel completeness.

**§7. Perpendiculars, midpoints and circles.** We have included two-point line-circle continuity in our axiom systems for ruler-and-compass geometry, since this corresponds to the the physical use of ruler and compass. Tarski, on the other hand, had segment-circle continuity. In this section, we will show how to construct perpendiculars and midpoints, using two-point line-circle continuity. We will deal with “dropped perpendiculars” (perpendiculars from a point  $p$  to a line  $L$ , assuming  $p$  is not on  $L$ ), and also with “erected perpendiculars” (perpendiculars to a line  $L$  at a point  $p$  on  $L$ ). Our results are of equal interest for classical and constructive geometry. As in [19], we abstain from the use of the dimension axioms or the parallel axiom (except in one subsection, where we mention our use of the parallel axiom explicitly); although these restrictions are not stated in the lemmas they are adhered to in the proofs.

We need to verify that the midpoint of a segment can be constructed by a term in intuitionistic Tarski geometry with Skolem functions. If we were willing to use the intersection points of circles, the problem might seem simple: we could use the two circles drawn in Euclid's Proposition I.1, and connect the two intersection points. (This is not Euclid's construction of midpoints, but still it is commonly used.) This matter is not as simple as it first appears, as we shall now explain.

We try to find the midpoint of segment  $pq$ . Let  $K$  be the circle with center  $p$  passing through  $q$ , and let  $C$  be the circle with center  $q$  passing through  $p$ , and let  $d$  and  $e$  be the two intersection points of these circles. Now the trick would be to prove that  $de$  meets  $pq$  in a point  $f$ ; if that could be done, then it is easy to prove  $f$  is the desired midpoint, by the congruence of triangles  $pef$  and  $qef$ . But it seems at first that the the full Pasch axiom is required to prove the existence of  $f$ . True, full Pasch follows from inner Pasch, at least classically, but we would have to verify that constructively using only the axioms of intuitionistic Tarski, which does not seem trivial. In particular, we will need the existence of midpoints of segments to do that!

In fact, the existence of midpoints has been the subject of much research, and it has been shown that one does not need circles and continuity at all! Gupta



Since circle-circle continuity enables us to construct an equilateral triangle on any segment (via Euclid I.1), we have justified the existence of midpoints and perpendiculars if circle-circle continuity is used. But that is insufficient for our purposes, since intuitionistic Tarski geometry does not contain circle-circle continuity as an axiom.

**7.2. A lemma of interest only constructively.** In erecting a perpendicular to line  $L$  at point  $a$ , we need to make use of a point not on  $L$  (which occurs as a parameter in the construction). In fact, we need a point  $c$  not on  $L$  such that  $ca$  is not perpendicular to  $L$ . Classically, we can make the case distinction whether  $ca$  is perpendicular to  $L$ , and if it is, there is “nothing to be proved”. But constructively, this case distinction is not allowed, so we must first construct such a point  $c$ . Our first lemma does that.

**LEMMA 7.2.** *Let point  $a$  lie on line  $L$ , and point  $s$  not lie on line  $L$ . Then there is a point  $c$  not on  $L$  such that  $ca$  is not perpendicular to  $L$ , and a point  $b$  on  $L$  such that  $\mathbf{B}(c, s, b)$ .*

*Remark.* Classically, this lemma is trivial, but constructively, there is something to prove.

*Proof.* Let  $b$  be a point on  $L$  such that  $ab = as$ . (Such a point can be constructed using only the segment extension axiom.) Then by Lemma 7.1,  $sb$  has a midpoint  $c$ , and  $ac \perp bc$ . We claim that  $ac$  cannot be perpendicular to  $L$ ; for it were, then triangle  $cab$  would have two right angles, one at  $c$  and one at  $a$ . That contradicts Satz 8.7. That completes the proof.

**7.3. Erected perpendiculars from triangle circumscription.** Following Szmielew, we have taken as our form of the parallel axiom, the axiom that given any three non-collinear points, there is another point equidistant from all three. (That point is then the center of a circumscribed circle containing the three given points.) An immediate corollary is the existence of midpoints and perpendiculars.

**LEMMA 7.3.** *Every segment has a midpoint and a perpendicular bisector. If  $p$  is a point not on line  $L$ , then there is a point  $x$  on  $L$  with  $px \perp L$ .*

*Proof.* Let  $ab$  be given with  $a \neq b$ . By Lemma 7.2, there is a point  $c$  such that  $a$ ,  $b$ , and  $c$  are not collinear. By triangle circumscription, there exists a point  $e$  such that  $ea = eb = ec$ . Then  $eab$  is an equilateral triangle. By Lemma 7.1,  $ab$  has a midpoint  $m$ . Since  $ea = eb$ , we have  $em \perp ab$ , by definition of perpendicular. That completes the proof.

While formally pleasing, there is something unsatisfactory here, because we intend that the axioms of our theories should correspond to ruler-and-compass constructions, and in order to construct the point required by the triangle circumscription axiom, we need to construct the perpendicular bisectors of  $ab$  and  $ac$ , and find the point of intersection (whose existence is the main point of the axiom). So from the point of view of ruler-and-compass constructions, our argument has been circular. Therefore the matter of perpendiculars cannot be left here. We must consider how to construct them with ruler and compass.

Once we have perpendiculars, several basic theorems follow easily. The following can be proved without using the parallel axiom or any continuity, from the assumption that dropped perpendiculars exist. Hence, without appealing to Gupta's construction, they can be proved in intuitionistic Tarski geometry (using triangle circumscription).

LEMMA 7.4. (i) *An exterior angle of a triangle is greater than either of the opposite interior angles.*

(ii) *The leg of a right triangle is less than the hypotenuse.*

(iii) *The triangle inequality holds.*

*Proof.* (i) is Satz 11.41 of [19]; (ii) is a special case of Satz 11.53. Neither of these proofs uses anything but elementary betweenness and congruence, and the existence of perpendiculars.

Turning to the triangle inequality, let triangle  $abc$  be given, and drop a perpendicular  $cx$  from  $c$  to  $\text{Line}(a, b)$ . If  $\mathbf{T}(a, x, b)$  then  $ab = ax + xb < ac + bc$  by (ii). Otherwise  $ab < ax < ac$ , so in either case we are finished. Hence constructively not not  $ab < ac + bc$ ; but segment inequality is stable, so  $ab < ac + bc$ . That completes the proof.

**7.4. Dropped perpendiculars from line-circle continuity.** In this section we discuss the following method (from Euclid I.12) of dropping a perpendicular from point  $p$  to line  $L$ : draw a large enough circle  $C$  with center  $p$  that some point of  $L$  is strictly inside  $C$ . Then apply two-point line-circle continuity to get two points  $u$  and  $v$  where  $C$  meets  $L$ . Then  $puv$  is an isosceles triangle, so it has a midpoint and perpendicular bisector. This argument is straightforwardly formalized in intuitionistic Tarski geometry:

LEMMA 7.5 (dropped perpendiculars from line-circle). *One can construct a perpendicular to line  $L$  from a point not on  $L$  by a ruler-and-compass construction, by essentially the construction of Euclid I.12, and prove the construction correct in intuitionistic Tarski geometry.*

*Proof.* Let  $p$  be a point, and let  $L$  be the line through two distinct points  $a$  and  $b$ . Suppose  $p$  is not on  $L$ . Let  $r = \text{ext}(p, a, a, b)$ , so that  $pr$  is longer than  $pa$ . Let  $C$  be the circle with center  $p$  passing through  $r$ . Then  $a$  is strictly inside  $C$ . By line-circle continuity, there are two points  $x$  and  $y$  on  $L$  where  $L$  meets  $C$ , i.e.  $px = py$ . Since by hypothesis  $p$  is not on  $L$ , the segment  $xy$  is the base of an isosceles triangle, so by Lemma 7.1 it has a midpoint  $m$ . Then  $pm \perp L$ , by definition of perpendicular. That completes the proof.

We note that the analogous lemmas with one-point line-circle or segment-circle in place of two-point line-circle will not be proved so easily. In the case of segment-circle, we would need to construct points outside  $C$  in order to apply the segment-circle axiom; but to show those points are indeed outside  $C$ , we would need the triangle inequality, which requires perpendiculars for its proof. In the case of one-point line-circle, we would need to construct the other intersection point, and the only apparent way to do that is to first have the dropped perpendicular we are trying to construct.

Of course, these difficulties are resolved if we are willing to use the 1965 discoveries of Gupta, who showed how to construct perpendiculars without any use

of circles. But that is beside the point here, since we are considering whether our choice of axioms corresponds well to Euclid or not. Also, the use of the triangle circumscription axiom is no help, since although we could prove the existence (if not the construction) of erected perpendiculars, we cannot do the same for dropped perpendiculars.

**7.5. Erected perpendiculars from dropped perpendiculars.** Next we prove the existence of erected perpendiculars, assuming only that we can drop perpendiculars, and without using any form of the parallel axiom. Gupta's proof, as presented in Satz 8.21 of [19], accomplished this. It is much simpler than Gupta's beautiful circle-free construction of dropped perpendiculars, but still fairly complicated. Gupta uses his Krippenlemma. Here we give a new proof, avoiding the use of the Krippenlemma, and using only simple ideas that might have been known to Tarski in 1959.

LEMMA 7.6 (erected perpendiculars). *In intuitionistic Tarski geometry with line-circle continuity but without circle-circle continuity, one can prove the following: Let  $L$  be a line; let  $a$  be a point on  $L$  and let  $s$  be a point not on  $L$ . Then there exists a point  $p$  on the opposite side of  $L$  from  $s$  such that  $pa \perp L$ . Moreover,  $pa$  is longer than the perpendicular from  $s$  to  $L$ .*

*The point  $r$  on  $L$  such that  $\mathbf{B}(p, r, s)$ , as well as the point  $p$ , are given by terms of intuitionistic Tarski geometry with Skolem functions.*



*Remarks.* No dimension axiom is used, and no parallel axiom. The point  $s$  and line  $L$  determine a plane in which  $p$  lies. We need  $s$  constructively also because we do not in general know how, without circle-circle continuity, to construct a point not on  $L$  by a uniform construction. We need  $pa$  longer than the perpendicular from  $s$  to  $L$  to avoid a case distinction in the construction of midpoints in the next lemma. "Shorter than" would be easier than "longer than", but we would need outer Pasch instead of inner Pasch, to prove that, and we don't have outer Pasch until we can erect perpendiculars.

*Proof.* By Lemma 7.2, we can find a point  $c$  not on  $L$ , and a point  $b \neq a$  on  $L$ , such that  $ca$  is not perpendicular to  $L$ , and  $\mathbf{B}(c, s, b)$ .

By Lemma 7.5, let  $x$  be a point on  $L$  such that  $cx \perp L$ . Then  $x \neq a$ , since  $ca$  is not perpendicular to  $L$ . Let  $d$  be the reflection of  $c$  in  $x$  and  $e$  the reflection of  $c$  in  $a$ . Since angle  $dxa$  is a right angle,  $da = ca$ . Since  $ea = ca$ , we have  $da = ea$ . Then triangle  $ade$  is isosceles, and hence has a midpoint by Lemma 7.1. Let  $p$  be that midpoint. Let  $y$  be the reflection of  $x$  in  $a$ , and  $f$  the reflection of  $d$  in  $a$ . Fig. 9 illustrates the situation.

Since reflection preserves congruence, we have  $xd = xc = ey$ , and  $yf = xd = xc = ey$ , and  $ca = da = fa$ . Since reflection preserves betweenness, we have  $\mathbf{B}(e, y, f)$ . (Thus  $f$  is the reflection of  $e$  in  $y$  as well as the reflection of  $d$  in  $a$ .) Let  $q$  be the reflection of  $p$  in  $a$ . Since  $ca = fa$ , angle  $aqf$  is a right angle. Therefore  $pc = pf$ . Now consider the four-point configurations  $(c, x, d, p)$  and  $(f, y, e, p)$ . Of the five corresponding pairs of segments, four are congruent, so by the 5-segment axiom,  $px = py$ . Since  $xa = ya$ , angle  $pax$  is a right angle and  $pa \perp L$ , as desired.

It remains to prove that  $p$  is on the other side of  $L$  from  $s$ . Fig. 10 illustrates the situation.

FIGURE 9. Erecting a perpendicular to  $L$  at  $a$ , given  $b$ ,  $s$ , and  $c$ .FIGURE 10.  $p$  is on the other side of  $L$  from  $s$ , by constructing first  $t$  and then  $r$ .

Segment  $xa$  connects two sides of triangle  $cde$ , and  $cp$  connects the vertex  $c$  to the third side, so by the crossbar theorem (Satz 3.17 of [19], derived with two applications of inner Pasch), there is a point  $t$  on  $xa$  (and hence on  $L$ ) with  $\mathbf{T}(c, t, p)$ . That is,  $p$  is on the other side of  $L$  from  $c$ . Using  $\mathbf{B}(c, s, b)$ , we can apply inner Pasch to the five points  $ptcbs$ , yielding a point  $r$  such that  $\mathbf{B}(t, r, b)$  and  $\mathbf{B}(p, r, s)$ . Since  $t$  and  $b$  lie on  $L$ , that shows that  $p$  and  $s$  are on opposite sides of  $L$ . That completes the proof.

**7.6. Midpoints from perpendiculars.** Next we intend to construct the midpoint of a non-degenerate segment  $ab$ . This is Satz 8.22 in [19]. That proof consists of a simple construction that goes back to Hilbert [13] (Theorem 26), and a complicated proof that it really constructs the midpoint. The proof of the correctness of Hilbert's construction from Tarski's axioms given in [19] is complicated, appealing to Gupta's "Krippenlemma", whose proof is not easy. Here we give another proof, not relying on the Krippenlemma. We do need to use the properties of reflection in a line. Since we can drop perpendiculars, we can define reflection in a line (for points not on the line). We need to know that reflection preserves betweenness and equidistance; otherwise put, reflection is an isometry. This is Satz 10.10 in [19], but the proof uses nothing but things

proved before the construction of perpendiculars and midpoints late in Chapter 8. Strangely, it is not explicitly stated in [19] that reflection in a line preserves betweenness, so we begin by proving that.

LEMMA 7.7. *Reflection in a line preserves betweenness (at least for points not on the line).*


*Proof.* Let  $\mathbf{B}(a, b, c)$ , and let  $p, q, r$  be the reflections of  $a, b, c$  in line  $L$ , and let  $u, v, w$  be the points on  $L$  at the feet of the perpendiculars from  $a, b, c$  to  $L$ . Suppose that  $q$  is not between  $p$  and  $r$ . Since reflection is an isometry, we have  $pq = ab$  and  $qr = bc$  and  $pr = ac$ . If  $q$  is not between  $p$  and  $r$ , then  $pqr$  is a triangle in which the sum of two sides  $pq$  and  $qr$  equals the third side  $pr$ . But dropping a perpendicular from  $q$  to  $pr$ , with foot at  $t$ , each of the two right triangles formed has its base less than its hypotenuse, so  $pr < pq + qr$ , contradiction. Hence  $\neg\neg\mathbf{B}(p, q, r)$ . By the stability of betweenness, we have  $\mathbf{B}(p, q, r)$ . That completes the proof.

LEMMA 7.8. *Let  $a \neq b$ , and suppose  $ap \perp ab$  and  $br \perp ab$ , and  $\mathbf{B}(a, m, b)$  and  $\mathbf{B}(p, m, r)$  and  $br = ap$ . Then  $m$  is the midpoint of  $ab$ , i.e.  $am = mb$ .*

*Proof.* By the stability of equality, we may use classical logic to prove  $am = mb$ ; hence we may refer to the proof in [19], page 65 (Abb. 31) (which appeals to the Krippenlemma, whose proof is complicated) without worrying whether it is constructive. But in the interest of giving a self-contained and simple proof, we will show how to finish the proof without appealing to the the complicated proofs of Gupta.

Let  $q$  and  $s$  be the reflections of  $p$  and  $r$ , respectively, in  $Line(a, b)$ . Fig. 11 illustrates the construction.

FIGURE 11. Given  $ap \perp ab$  and  $br \perp ab$  and  $br = ap$ , and  $ab$  meets  $pr$  at  $m$ , then  $m$  is the midpoint of  $ab$ .




Since  $ap \perp ab$  and  $br \perp ab$  we have  $\mathbf{B}(q, a, p)$  and  $\mathbf{B}(r, b, s)$ . Since reflection in a line preserves betweenness, and  $m$  is its own reflection since  $\mathbf{B}(a, m, b)$ , we have  $\mathbf{B}(q, m, s)$ . Hence  $m$  is the intersection point of the diagonals of the

quadrilateral  $qpsr$ . Since reflection in a line is an isometry, we have  $qr = ps$ . Since  $qa = ap = rb = bs$ , we have  $qp = rs$ . Hence the opposite sides of quadrilateral  $qpsr$  are equal. By Satz 7.29, the diagonals bisect each other. Hence  $mp = mr$  and  $mq = ms$ . Now by the inner five-segment theorem (Satz 4.2), applied to the configurations  $qapm$  and  $sbrm$ , we have  $ma = mb$ . Then  $m$  is the midpoint of  $ab$ . That completes the proof.

LEMMA 7.9 (midpoint existence). *In intuitionistic Tarski geometry with only line-circle continuity, midpoints exist. More precisely, given segment  $ab$  (with  $a \neq b$ ) and point  $s$  not collinear with  $ab$ , one can construct the midpoint  $m$  of  $ab$ .*

*Remark.* The proof shows how to derive the existence of midpoints from the existence of (erected) perpendiculars; we have shown above that line-circle continuity enables us to erect perpendiculars.

FIGURE 12. Midpoint from erected perpendiculars



*Proof.* The construction is illustrated in Fig. 12. Let segment  $ab$  be given along with point  $s$  not on  $L = \text{Line}(a, b)$ . Erect a perpendicular  $qb$  to  $ab$  at  $b$  (on the opposite side of  $L$  from  $s$ ). Then erect a perpendicular  $ap$  to  $L$  at  $a$ , with  $p$  on the opposite side of  $L$  from  $q$ , and nearer to  $L$  than  $q$ , by Lemma 7.6. Since  $p$  is on the opposite side of  $L$  from  $q$ , there is a point  $t$  on  $L$  with  $\mathbf{B}(p, t, q)$ . Then construct point  $r$  on segment  $qb$  so that  $rb = ap$ .

Applying inner Pasch to the five-point configuration  $ptqrb$ , we find a point  $m$  such that  $\mathbf{B}(a, m, b)$  and  $\mathbf{B}(p, m, r)$ . Note that by means of the extra condition in Lemma 7.6, we have guaranteed that a uniform construction (given by a single term of intuitionistic Tarski with Skolem functions) constructs the midpoint. By Lemma 7.8,  $m$  is the midpoint of  $ab$ . That completes the proof.

LEMMA 7.10. *The midpoint of segment  $ab$  is given by a term of intuitionistic Tarski geometry with Skolem functions,  $\text{midpoint}(a, b, s)$ , assuming that  $a \neq b$  and  $s$  does not lie on  $\text{Line}(a, b)$ .*

*Remark.* Of course since the midpoint is unique, the value does not depend on the parameter  $s$ . Nevertheless we do not know how to get rid of  $s$  in the term, as we need a point not on  $Line(a, b)$  to construct a perpendicular to  $ab$ .

*Proof.* Hilbert's construction is given by the following script, in which  $Perp(a, b, s)$  and  $wit(a, b, s)$  are the terms giving the construction in Lemma 7.6; that is, the first is a point on the perpendicular and the second is a witness that it is on the other side of  $Line(a, b)$ .

```

midpoint(a, b, s){
  p = Perp(a, b, s)    Then ap ⊥ ab
  q = Perp(b, a, p)    Then bq ⊥ ab on the other side of ab from p
  t = ext(q, b, q, b)  The reflection of r in b
  r = ext(t, q, a, p)  Then ap = br.
  x = wit(a, b, s)     Then B(a, x, b) and B(p, x, q) and T(b, r, q)
  M = ip(p, x, q, b, r) Then T(x, M, b) and T(a, M, b)
  return M }

```

**COROLLARY 7.11.** *In intuitionistic Tarski geometry (with two-point line-circle but without any parallel axiom) we can construct both dropped and erected perpendiculars, and midpoints.*

*Proof.* We have justified Euclid I.12 for dropped perpendiculars, and shown how to construct erected perpendiculars from dropped perpendiculars, and midpoints from erected perpendiculars. That completes the proof.

That result is pleasing, but it leaves open the question of whether the axiom system required can be weakened. It is not obvious how to weaken it even to one-point line-circle or segment-circle continuity; but Gupta showed in 1965 that no continuity whatsoever is required. We discuss his construction in the next section.

**7.7. Gupta's perpendicular construction.** Here we give Gupta's beautiful construction of a dropped perpendicular. It can be found in his 1965 thesis [11] and again in [19], p. 61.

The initial data are two distinct points  $a$  and  $b$ , and a third point  $c$  not collinear with  $ab$ . The construction goes as follows: Extend  $ba$  by  $ac$  to produce point  $y$ . Then  $acy$  is an isosceles triangle, so by Lemma 7.1, we can construct its midpoint  $p$ , and  $apc$  is a right angle. Now extend segment  $cy$  by  $ac$  to point  $q$ , and extend  $ay$  by  $py$  to produce point  $z$ . Then reflect  $q$  in  $z$  to produce  $q'$ . Then extend  $q'y$  by  $yc$  to produce point  $c'$ . By construction  $cy c'$  is an isosceles triangle, so its base  $cc'$  has a midpoint  $x$ , which is the final result of the construction. It is not immediately apparent either that  $x$  is collinear with  $ab$  or that  $cx \perp ax$ , but both can be proved. The proof occupies a couple of pages, but it uses only (A1)-(A7); in other words, no continuity, no dimension axioms, no parallel axiom. By the stability of equality, if  $cx \perp ax$  can be classically proved, it can be constructively



The construction starts with a line  $L$ , a point  $a$  on  $L$ , and a point  $c$  not on  $L$  such that  $ca$  is not perpendicular to  $L$ . Then we draw the circle with center  $c$  through  $a$ , and let  $b$  be the other point of intersection with  $L$ . Then the line  $bc$  meets the circle at a point  $e$ , and  $ea$  is the desired perpendicular to  $L$  (although we have not proved it here).

The correctness of this construction, i.e. that  $ea$  is indeed perpendicular to  $L$ , is equivalent to Euclid III.31, which says an angle inscribed in a semicircle is a right angle. Euclid's proof uses I.29, which in turn depends on I.11, which requires erecting a perpendicular. Since III.31 is certainly not valid in hyperbolic geometry, we will need to use the parallel postulate in any proof of III.31, and it seems extremely unlikely that we can prove III.31 without first having proved the existence of erected perpendiculars. Thus, this construction cannot replace the ones discussed above in the systematic development of geometry. Still, it is of interest because it is the shortest possible ruler-and-compass construction.

**§8. Other forms of the parallel axiom.** Within neutral geometry (that is, geometry without any form of the parallel postulate), we can consider the logical relations between various forms of the parallel axiom. In [5], we considered the Playfair axiom, Euclid 5, and the strong parallel axiom, which are all classically equivalent to Euclid 5. Constructively, Playfair is weaker, as shown in [5]; a formal independence result confirms the intuition that it should be weaker because it makes no existential assertion. The other versions of the parallel postulate, which do make existential assertions, each turn out to be fairly easy to prove equivalent to either Euclid 5 or the strong parallel postulate. In [5], we prove that Euclid 5 and the strong parallel postulate are actually constructively equivalent, although the proof requires first developing the geometrical definitions of arithmetic using only Euclid 5.


In [5], we gave the proof that the triangle circumscription principle is equivalent to the strong parallel axiom; below we prove that Tarski's version of the parallel postulate taken as axiom (A10) in [19], is equivalent to Euclid 5. Lemmas in this section are proved in neutral geometry, i.e. without any form of the parallel postulate. It follows that all the known versions of the parallel postulate (that are equivalent in classical Tarski geometry with line-circle continuity) that make an existential assertion are also equivalent in constructive Tarski geometry, although some of the proofs are much longer.

LEMMA 8.1. *Playfair's axiom implies the alternate interior angle theorem, that any line traversing a pair of parallel lines makes alternate interior angles equal.*

*Proof.* Since ordering of angles is stable, we can argue by contradiction. Hence the usual classical proof of the theorem applies.

**8.1. Euclid 5 formulated in Tarski's language.** Here we give a formulation of Euclid's parallel postulate, expressed in Tarski's points-only language. Euclid's version mentions angles, and the concept of "corresponding interior angles" made by a transversal. The following is a points-only version of Euclid 5. See Fig. 15.


FIGURE 15. Euclid 5. Transversal  $pq$  of lines  $M$  and  $L$  makes corresponding interior angles less than two right angles, as witnessed by  $a$ . The shaded triangles are assumed congruent. Then  $M$  meets  $L$  as indicated by the open circle.



$$\mathbf{B}(q, a, r) \wedge \mathbf{B}(p, t, q) \wedge pr = qs \wedge pt = qt \wedge rt = st \wedge \neg Col(s, q, p) \quad (\text{Euclid 5}) \\ \wedge s \neq q \rightarrow \exists x \mathbf{B}(p, a, x) \wedge \mathbf{B}(s, q, x)$$

**8.2. Tarski's parallel axiom.** As mentioned above, Tarski [24] and later [19] took a different form of the parallel postulate, illustrated in Fig. 16.<sup>16</sup> The following axiom is similar to Tarski's axiom, and we give it his name, but his axiom used non-strict betweenness and did not include the hypothesis that  $a$ ,  $b$ , and  $c$  are not collinear. It is intended to say that if  $t$  is in the interior of angle  $bac$ , then there is a line through  $t$  that meets both sides of the angle. To express this using variables for points only, Tarski used the point  $d$  to witness that  $t$  is in the interior of the angle. See Fig. 16.

FIGURE 16. Tarski's parallel axiom



The degenerate cases are trivial: if  $a$ ,  $b$ , and  $c$  are collinear, then we can (classically, or with more work also constructively) find  $x$  and  $y$  without any parallel axiom, and if (say)  $d = b$  then we can take  $x = t$  and  $y = c$ , etc. Hence the following axiom is classically equivalent in neutral geometry to the one used by Tarski:

<sup>16</sup>Technically, according to [25], the axioms taken *op. cit* differed in the order of arguments to the last betweenness statement, but that is of no consequence.


$$\begin{aligned}
 & \mathbf{B}(a, d, t) \wedge \mathbf{B}(b, d, c) \wedge a \neq d && \text{(Tarski parallel axiom)} \\
 & \wedge (\neg \mathbf{B}(a, b, c) \wedge \neg \mathbf{B}(b, c, a) \wedge \neg \mathbf{B}(c, a, b)) \\
 & \rightarrow \exists x \exists y (\mathbf{B}(a, b, x) \wedge \mathbf{B}(a, c, y) \wedge \mathbf{B}(x, t, y))
 \end{aligned}$$

Something like this axiom was first considered by Legendre (see [10], p. 223), but he required angle  $bac$  to be acute, so Legendre's axiom is not exactly the same as Tarski's parallel axiom. The axiom says a bit more than just that  $xy$  meets both sides of the angle, because of the betweenness conditions in the conclusion; but it would be equivalent to demand just that  $x$  and  $y$  lie on the rays forming angle  $bac$ , as can be shown using Pasch.

**8.3. Euclid 5 implies Tarski's parallel axiom.**

**THEOREM 8.2.** *Euclid 5 implies Tarski's parallel axiom in neutral intuitionistic Tarski geometry.*

FIGURE 17. Constructive proof of Tarski's parallel axiom from Euclid 5.  $M$  is constructed parallel to  $Line(b, c)$  and  $cd = ce$  and  $bd = bg$ . Then  $x$  and  $y$  exist by Euclid 5.



*Proof.* Assume the hypothesis of Tarski's parallel axiom. Construct line  $M$  parallel to  $Line(b, c)$  through  $t$ . Construct point  $e$  by extending segment  $dc$  by  $dc$ ; then  $ec = dc$  and  $\mathbf{B}(d, c, e)$ , as illustrated in Fig. 17. Let  $L$  be  $Line(a, c)$ . Then  $Line(d, t)$  meets  $L$  at  $a$ , and does not coincide with  $L$  since, if it did coincide with  $L$ , then points  $d$  and  $c$  would be on  $L$ , and hence point  $b$ , which is on  $Line(b, c)$ , would lie on  $L$  by Axiom I3; but that would contradict the hypothesis that  $a$ ,  $b$ , and  $c$  are not collinear. Hence  $Line(d, t)$  meets  $L$  only at  $a$ , by Axiom I3. Hence segment  $dt$  does not meet  $L$ . By outer Pasch (applied to  $adtec$ ), there is a point  $f$  on  $L$  with  $\mathbf{B}(e, f, t)$ . Now we apply Euclid 5; the two parallel lines are  $Line(b, c)$  and  $M$ , and the conclusion is that  $L$  meets  $M$  in some point, which we call  $y$ . Specifically we match the variables  $(L, K, M, p, r, a, q)$  in Euclid 5 to the following terms in the present situation:  $(M, Line(b, c), Line(a, c), c, e, f, t)$ . Then all the hypotheses have been proved, except that we have  $\mathbf{B}(e, f, t)$  while



$\mathbf{B}(p, v, q)$  and  $\mathbf{B}(u, v, a)$ . (This is where we use the hypothesis  $\mathbf{B}(r, a, q)$ .) Then  $v$  witnesses that  $t$  is in the interior of angle  $upa$ . By Tarski's parallel axiom, there exist points  $x$  and  $y$  with  $\mathbf{B}(x, u, p)$  and  $\mathbf{B}(p, a, y)$ . Then line  $L$  meets side  $xy$  of triangle  $xpy$  at  $q$ , and does not meet the closed side  $xp$ , since  $K$  is parallel to  $L$ ; so by Pasch's theorem (Corollary 10.8), line  $L$  meets segment  $py$  in a point  $e$ .

We next wish to prove  $\mathbf{B}(p, a, e)$ . By the stability of betweenness, we may prove it by contradiction. By hypothesis,  $r$  does not lie on  $L$ , so  $a$  does not lie on  $L$ . Hence  $a \neq e$ . Since  $L$  and  $K$  are parallel,  $e \neq p$ . Suppose  $\mathbf{B}(p, e, a)$ . Then  $L$  meets side  $pa$  of triangle  $pra$ .  $L$  does not meet the closed side  $pr$  since  $L$  is parallel to  $K$ . Therefore by Pasch's theorem,  $L$  meets the open segment  $qr$ . But then  $L$  contains two distinct points of  $\text{Line}(q, r)$ , so  $a$  lies on  $L$ , contradiction. Hence  $\neg\mathbf{B}(p, e, a)$ . The only remaining possibility is  $\mathbf{B}(e, p, a)$ . In that case,  $e$  is on the opposite side of  $K$  from  $a$ . But  $aq$  does not meet  $K$ , since if it did,  $\text{Line}(q, r)$  would coincide with  $K$ , but  $a$  does not lie on  $K$ . That is,  $q$  and  $a$  are on the same side of  $K$ . By the plane separation theorem (Theorem 2.5)  $q$  and  $e$  are on the opposite side of  $K$ . Hence there is a point of  $K$  on  $qe$ , contradicting the fact that  $K$  and  $L$  are parallel. That completes the proof of  $\mathbf{B}(p, a, e)$ .

We still must show  $\mathbf{B}(s, q, e)$ . Since we now have  $\mathbf{B}(p, a, e)$ , we can apply outer Pasch to  $paesq$  to conclude  $\mathbf{B}(s, q, e)$ . That completes the proof of the theorem.

**§9. Uniform perpendicular and uniform rotation.** In classical geometry there are two different constructions, one for “dropping a perpendicular” to line  $L$  from a point  $p$  not on  $L$ , and the other for “erecting a perpendicular” to  $L$  at a point  $p$  on  $L$ . A “uniform perpendicular” construction is a method of constructing, given three points  $a, b$ , and  $x$ , with  $a \neq b$ , a line perpendicular to  $\text{Line}(a, b)$  and passing through  $x$ , without a case distinction as to whether  $x$  lies on  $L$  or not.


In constructive geometry, it is not sufficient to have dropped and erected perpendiculars; we need uniform perpendiculars. Similarly, we need uniform rotations: to rotate a given point  $x$  on  $\text{Line}(c, a)$  clockwise about center  $c$  until it lies on a given line through  $c$ , without a case distinction as to whether  $\mathbf{B}(x, c, a)$  or  $x = c$  or  $\mathbf{B}(c, x, a)$ . We also need uniform reflections: to be able to reflect a point  $x$  in a line  $L$  without a case distinction whether  $x$  is on  $L$  or not. It will turn out (not surprisingly) that the three problems of perpendiculars, rotations, and reflections are closely related.

**9.1. Uniform perpendicular using line-circle.** In this section, we take up the construction of the uniform perpendicular.

**THEOREM 9.1.** *Uniform perpendiculars can be constructed in intuitionistic Tarski geometry, using two-point line-circle continuity.*

*Proof.* The idea is simple: Draw a circle  $C$  around  $x$  whose radius exceeds  $ax$ . Then by two-point line-circle continuity there are distinct points  $p$  and  $q$  on  $L = \text{Line}(a, b)$  that lie on  $C$ . Then the perpendicular bisector of segment  $pq$  is the desired perpendicular. The matter is, however, trickier than the similar verification of Euclid I.2, of the requirement to find a point  $r$  through which to draw the circle  $C$ . We must have  $xr > ax$ , but we are not allowed to make a

FIGURE 19. The simplest uniform perpendicular construction.  $M = \text{Perp}(x, L)$  is constructed perpendicular to  $L$  without a case distinction whether  $x$  is on  $L$  or not. Draw a large enough circle  $C$  about  $x$ . Then bisect  $pq$  and erect  $K$  at the midpoint. The radius is  $ac = ab + ax$ .



case distinction whether  $x = a$  or not, and since we are not allowed to extend a null segment, we cannot find  $r$  by extending the (possibly null) segment  $ax$ . First we construct point  $c$  by extending segment  $ab$  by  $ax$ . Since  $ab$  is not null, and we are allowed to extend a non-null segment by a possibly null segment, this is legal. Then we make use of the construction  $e_2$  corresponding to Euclid I.2, developed in Lemma 11.6. We define  $r = e_2(x, a, c)$ . Then  $ar = ac$  and since  $ac > ax$  we have  $ar > ax$ . Now draw the circle  $C$  with center  $x$  through  $r$ . Then  $a$  is inside  $C$ , so  $p$  and  $q$  exist by two-point line-circle continuity, and we can complete the proof using any construction of the perpendicular bisector, for example the one developed using two-point line circle continuity only, or Gupta's more complicated one.

It is not difficult to construct a script, and hence in principle a term, describing this construction. This term does require an "extra" parameter  $s$ , for a point assumed to be not on the line  $L$ , in order to erect the required perpendicular. The script is sketched in the caption of the figure.

**9.2. Uniform perpendicular without any continuity.** The above construction has one disadvantage: it relies on line-circle continuity. Although our main interest is in intuitionistic Tarski geometry, nevertheless it is of some interest to study the theory that results from deleting line-circle continuity as an axiom, i.e. the intuitionistic counterpart of Tarski's (A1)-(A10). It turns out, one can also construct a uniform perpendicular in that theory, although of course, one must use Gupta's perpendiculars. It turns out that we also need the parallel axiom to construct uniform perpendiculars, although it is not needed for Gupta's perpendiculars.

We need the following lemmas.<sup>18</sup>

LEMMA 9.2. *Let  $abcd$  be a quadrilateral with two adjacent right angles at  $a$  and  $d$ , and two opposite sides equal, namely  $ab = cd$ . Suppose  $a$  and  $d$  are on the same side of  $bc$ . Then also the other pair of opposite sides are equal and  $abcd$  is a rectangle.*

*Remarks.* The proof necessarily will require the parallel axiom, as the lemma is false in hyperbolic geometry. The assumption that  $a$  and  $d$  are on the same side of  $bc$  ensures that the quadrilateral lies in a plane (we do not use any dimension axiom in the proof).

*Proof.* We first claim  $ad$  and  $bc$  are parallel. By definition, two lines are parallel if they lie in a plane and do not meet, so it suffices to show  $ad$  and  $bc$  do not meet. Suppose they meet at point  $p$ . Let  $J$  be the perpendicular bisector of  $ad$ ; then since  $ab = cd$ , reflection in  $J$  takes  $ab$  onto  $cd$ , and hence fixes  $\text{Line}(b, c)$  and  $\text{Line}(a, d)$ . Hence  $ad$  and  $bc$  also meet at the reflection  $q$  of  $p$  in  $J$ . But distinct lines cannot meet in two points; hence  $p$  is on  $J$ , i.e. between  $a$  and  $d$ . Hence  $a$  and  $d$  are on opposite sides of  $bc$ . But by hypothesis they are on the same side of  $bc$ , contradicting Satz 9.9 of [19]. That proves  $ad$  and  $bc$  are parallel. Also  $ab$  and  $cd$  are parallel, since if they meet at a point  $u$  then there would be two perpendiculars from  $u$  to  $\text{Line}(a, b)$ , contradicting the first part of Satz 8.18. Let  $m$  be the midpoint of  $bc$ , and let  $e$  be the reflection of  $d$  in  $m$ . By Euclid 5,  $eb$  is parallel to  $ad$ , since triangle  $emb$  is congruent to triangle  $dma$  (by reflection in  $m$ ). But it is also a consequence of the parallel axiom that the parallel to  $ad$  through  $b$  is unique. Hence  $e$  lies on  $\text{Line}(b, c)$ . Since  $b$  is the reflection of  $a$  in  $m$ ,  $\text{Line}(b, c)$  is the reflection of  $\text{Line}(a, d)$  in  $m$ , and consequently the angles at  $b$  and  $c$  are right angles. Now triangles  $abc$  and  $cda$  are congruent, since they are right triangles with one congruent leg and a common hypotenuse. Then the other legs are congruent. But that is the desired conclusion,  $ad = bc$ . That completes the proof.


LEMMA 9.3. *(in intuitionistic Tarski geometry without continuity) Let  $abcd$  be a quadrilateral whose diagonals  $ac$  and  $bd$  bisect each other at  $x$ . Then (i) opposite sides of  $abcd$  are parallel, and (ii) the lines connecting midpoints of opposite sides pass through  $x$ , and (iii) they are parallel to the other sides, as shown in Fig. 20.*

*Proof.* Since reflection (in point  $x$ ) preserves congruence (Satz 10.10 of [19]), we have  $ab = cd$  and  $bc = ad$ , i.e. opposite sides are equal. Let  $m$  be the midpoint of  $ab$  and  $k$  the reflection of  $m$  in  $x$ . Since reflection preserves congruence,  $ck = kd$ , so  $k$  is the midpoint of  $cd$ . Let  $n$  be the midpoint of  $bc$  and  $\ell$  the midpoint of  $ad$ . Then similarly  $n\ell$  passes through  $x$ . That proves that the lines connecting midpoints of opposite sides pass through  $x$  as claimed in (ii) of the theorem.

---

<sup>18</sup>Though these are fairly routine exercises in Euclidean geometry, we need to verify that they are provable in intuitionistic Tarski geometry without any continuity. These theorems do not occur in [19], and even if they had occurred, we would still need to check that they are provable with triangle circumscription instead of Tarski's parallel axiom. In fact we will use Euclid 5, which is provable from triangle circumscription, as shown in [5].

FIGURE 20. Given  $ac$  and  $bd$  bisect each other at  $x$ . Then the lines that look parallel, are parallel.



Suppose point  $u$  lies on  $Line(a, d)$  and also on  $Line(b, c)$ . Let  $v$  be the reflection of  $u$  in  $x$ . Then since reflection preserves collinearity, by Lemma 7.7,  $v$  also lies on both lines. But  $v \neq u$ , since if  $v = u$  then  $v = x = u$ , but  $m$  does not lie on  $Line(a, d)$ . Hence there are two distinct points  $u$  and  $v$  on  $Line(a, d)$  and  $Line(b, c)$ , contradiction. Hence those two lines are parallel, as claimed. Similarly, the other two sides of the quadrilateral are parallel. That proves part (i) of the theorem.

By the inner five-segment theorem (Satz 4.2) applied to  $ambx$  and  $ckdx$ , we have  $mx = xk$ . Therefore triangle  $ncx$  is congruent to triangle  $lax$ , since they have all three pairs of corresponding sides equal. Similarly triangle  $ckx$  is congruent to triangle  $amx$ .


We next claim that  $mk = ad$ . By the stability of equality, we may argue by contradiction and cases. Drop a perpendicular from  $d$  to  $mk$ ; let  $f$  be the foot. Case 1,  $f = k$ . Then  $dk \perp mk$ , and by reflection in  $x$ ,  $bm \perp mx$ , so  $ma \perp mk$  and  $mkda$  has two adjacent right angles. Hence its opposite sides  $mk$  and  $ad$  are equal, by Lemma 9.2, since  $a$  and  $d$  are on the same side of  $mk$  because  $ab$  and  $bd$  both meet  $Line(m, k)$  (in  $m$  and  $x$  respectively). Case 2,  $\mathbf{B}(m, f, k)$ . Then let  $g$  be the reflection of  $f$  in  $x$ , and  $h$  the reflection of  $g$  in  $m$ . Then the triangles  $dfk$ ,  $bgm$ , and  $ahm$  are congruent right triangles (by reflection), and  $\mathbf{B}(h, m, g)$ . Then  $hm = mg = fk$ . Since  $hadf$  has two adjacent right angles, its opposite sides  $hf$  and  $ad$  are equal, since again  $a$  and  $d$  are on the same side of  $Line(h, f)$ , so Lemma 9.2 applies. But  $hf = mk$ , since they differ by adjoining and removing equal segments  $hm$  and  $fk$ . Hence  $mk = hf = ad$ , so  $mk = ad$  as claimed. Case 3,  $\mathbf{B}(m, k, f)$ . Reflection in the line  $mk$  reduces this case to Case 2. That completes the proof that  $mk = ad$ .

Now  $mkda$  is a parallelogram with opposite sides equal. Therefore, part (i) of the theorem, which has already been proved, can be applied to it. Hence  $mk$  is parallel to  $ad$  as claimed. That completes the proof.

LEMMA 9.4. (*in intuitionistic Tarski geometry without continuity*) Let line  $J$  be parallel to line  $M$ , and suppose  $M \perp L$ . Then  $J$  meets  $L$  in a point  $x$  and  $J$  is perpendicular to  $L$  at  $x$ .

*Proof.* See Fig. 21. Let  $p$  be the point of intersection of  $M$  and  $L$ . Drop a perpendicular from  $p$  to  $J$ ; let  $x$  be the foot of that perpendicular. Let  $m$  be the midpoint of  $px$  (using Gupta's midpoint to avoid any need for continuity). Let  $q$  be any point other than  $p$  on  $M$  and let  $r$  be the reflection of  $q$  in  $p$ . Then  $mq = mr$  since  $M \perp L$ . Let  $t$  and  $s$  be the reflections of  $q$  and  $r$  in  $m$ ,

FIGURE 21. Given  $M \perp L$  and  $J$  parallel to  $M$ , show  $J \perp L$  and construct the intersection point  $x$  of  $J$  and  $L$  by dropping a perpendicular from  $p$  to  $J$ . Construct  $r$ ,  $t$ , and  $s$  by reflection. Then  $ts$  is parallel to  $M$  and hence lies on  $J$ , and  $ts \perp L$ .



respectively. Since reflection preserves congruence,  $sx = pq$ . Then  $xs$  is parallel to  $M$ , since if point  $u$  on  $M$  lies on  $Line(x, s)$ , then the reflection of  $u$  in  $m$  is another point on both lines, contradiction. Now  $J$  is by hypothesis parallel to  $M$ , and  $J$  passes through  $x$ . But by the parallel axiom, there cannot be two parallels to  $M$  through  $x$ . Hence the two coincide:  $J = Line(x, s)$ . We have  $ms = mq$  by reflection,  $mq = mr$  since  $M \perp L$ , and  $mr = mt$  by reflection; hence  $ms = mt$ . Then  $J \perp L$  by the definition of perpendicular. That completes the proof.


**THEOREM 9.5.** [*Uniform perpendicular*] *Uniform perpendiculars can be constructed in intuitionistic Tarski geometry without any continuity axioms. Explicitly: there is a term  $Project(x, a, b, w)$  in intuitionistic Tarski with Skolem functions, such that if  $a \neq b$  and  $w$  is any point not on  $L = Line(a, b)$ , and  $f = Project(x, a, b, w)$ , then  $Col(a, b, f)$ , and the erected perpendicular to  $L$  at  $f$  contains  $x$ .*

*Remark.* The two main points of the lemma are that we do not need a case distinction whether  $x$  is on  $L$  or not, and we do not use any continuity axiom. But we do use the parallel axiom.

*Proof.* See Fig. 22. We begin by constructing a point  $p$  on  $L$  such that  $px$  is not perpendicular to  $L$ . (Of course that is trivial classically, but constructively, there is something to prove. ) The line  $L$  is given by two points  $a$  and  $b$ ; let  $p$  be the result of extending  $ab$  by  $ab$  to point  $q$  and then extending  $aq$  by  $xb$  to point  $p$ . Then  $\mathbf{B}(b, q, p)$  and  $qp = xb$ . We claim  $x$  lies on  $L$ . By the stability of incidence, we may prove that by contradiction. If  $x$  does not lie on  $L$ , then  $bpq$  is a triangle, and it has leg  $bp$  greater than the hypotenuse  $bx = qp$ , contradiction. Therefore  $x$  is on  $L$  as claimed. Then  $px$  lies on  $L$  too, and  $p \neq x$ ; hence  $px$  is not perpendicular to  $x$ . That completes the preliminary construction of  $p$ .

Erect a perpendicular  $qp$  to  $L$  at  $p$ . (For that we need the point  $w$  not on  $L$ .) Let  $s$  be the reflection of  $q$  in  $x$ . Let  $t$  be the reflection of  $p$  in  $x$ . Then  $tx = xp$  and  $sx = xq$ . By Lemma 9.3,  $stqp$  is a parallelogram whose diagonals bisect each other. Let  $m$  be the midpoint of  $tq$  and  $k$  the midpoint of  $sp$ . By

FIGURE 22. Uniform perpendicular. Given  $L$  and  $x$  construct perpendicular  $J$  to  $L$  passing through  $x$ , without a case distinction whether  $x$  is on  $L$  or not.



Lemma 9.3,  $tq$  is parallel to  $sp$ , so  $m \neq k$ . We claim  $J = \text{Line}(k, m)$  is the desired perpendicular to  $L$ . By Lemma 9.3,  $J$  is parallel to  $qp$ . But  $qp \perp L$  by construction. By the parallel axiom, lines parallel to  $qp$  are also perpendicular to  $L$ ; so  $J \perp L$ . By Lemma 9.4, the intersection point  $f$  of  $J$  and  $L$  exists and  $J \perp L$ . The point  $f$  is the value of  $\text{Project}(x, a, b, s)$ . That completes the proof.

*Remark.* We do not know how to construct uniform perpendiculars without using either the parallel axiom or two-point line-circle, although either one suffices, and any form of the parallel axiom suffices (because we just need a few simple lemmas about parallelograms).

**9.3. Uniform Reflection.** Another construction from [5] that we need to check can be done with Tarski's Skolem symbols is the "uniform reflection". The construction  $\text{Reflect}(x, a, b, s)$  gives the reflection of point  $x$  in  $L = \text{Line}(a, b)$ , without a case distinction as to whether  $x$  is on  $L$  or not. (The parameter  $s$  is some point not on  $L$ .) First we note a difficulty: even though we can define  $f = \text{Project}(x, a, b, s)$ , we cannot just extend segment  $xf$  by  $xf$ , since  $xf$  might be a null segment, and in constructive geometry, we can only extend non-null segments.

The solution given in [5] is to first define rotations, and then use the fact that the reflection of  $x$  in  $\text{Line}(a, b)$  is the same as the result of two ninety-degree rotations of  $x$  about  $f = \text{Project}(x, a, b, s)$ . The construction given for rotations in [5] only involves bisecting the angle in question, and dropping two perpendiculars to the sides, none of which is problematic in Tarski's theory.

#### 9.4. Equivalence of line-circle and segment-circle continuity.

LEMMA 9.6. *Two-point line-circle, one-point line-circle, and segment-circle continuity are equivalent in (A1)-(A10).*

*Remark.* This proof depends heavily on Gupta's 1965 dissertation.

*Proof.* Because of Gupta, we have dropped perpendiculars, and we have shown above that from dropped perpendiculars, erected perpendiculars, midpoints, and uniform perpendiculars follow. Then one-point line-circle implies two-point line-circle, by reflection in the uniform perpendicular from line  $L$  through the center of the circle. Note that classically, the case when  $L$  passes through the center is trivial, so a dropped perpendicular is enough, and the parallel axiom is not required.

Two-point line-circle implies segment-circle immediately.

Next we will show that segment-circle implies one-point line-circle. It suffices to construct a point  $q$  on  $L$  outside the circle. Let  $z$  be any point on  $L$  different from  $p$ , and define  $q$  by extending segment  $zp$  by three times the radius  $ab$ . Then  $qp > ab - ap \geq qp - ab \geq 2ab > ab$ , by the triangle inequality. (This short piece of algebra can be converted to a lengthier, purely geometric, proof.) By the stability of betweenness, it suffices to prove this classically, so we may divide into cases as to whether  $L$  is a diameter or not; if it is, we are finished, so we may assume it is not. Then we have a non-degenerate triangle. The triangle inequality for a non-degenerate triangle is proved by dropping a perpendicular from the vertex to the base, and using the theorem that the leg of a right triangle is less than the hypotenuse (Lemma 7.4). That completes the proof.

*Remark.* One may wonder why Tarski chose segment-circle rather than line-circle continuity as an axiom. It might be because segment-circle continuity asserts the existence of something that turns out to be unique; but that consideration did not bother Tarski when he chose (A10) as his parallel axiom. More likely it was just shorter. Although the diagram for (either form of) line-circle continuity is simpler, the formal expression as an axiom is longer, especially if collinearity is written out rather than abbreviated; and Tarski placed importance on the fact that his axioms could be written intelligibly without abbreviations.

**9.5. Representation theorems.** The following important theorem was stated in 1959 by Tarski [24]:

**THEOREM 9.7.** (i) *The models of classical Tarski geometry with segment-circle continuity are the planes  $F^2$  over a Euclidean field  $F$ .*

(ii) *The models of classical Tarski geometry with no continuity axioms are the planes  $F^2$  over a Pythagorean field  $F$ .*<sup>19</sup>

*Remark.* Tarski wrote "ordered field" instead of "Pythagorean field" in (ii), but one needs to be able to take  $\sqrt{1+x^2}$  in  $\mathbb{F}$  to verify the segment extension axiom in  $\mathbb{F}^2$ , as Tarski surely knew.

There is also a version of the representation theorem for intuitionistic Tarski geometry. For that to make sense, we must define ordered fields and Euclidean fields constructively. That is done in [5], as follows: We take as axioms the stability of equality and "Markov's principle" that  $\neg x \leq 0$  implies  $x > 0$ , which is

---

<sup>19</sup>A Pythagorean field is one in which sums of two squares always have square roots, or equivalently,  $\sqrt{1+x^2}$  always exists, as opposed to Euclidean in which all positive elements have square roots.

similar to the stability of betweenness in geometry. Then we require that nonzero elements have multiplicative inverses; just as classically, a Euclidean field is an ordered field in which positive elements have square roots (and a Pythagorean field is one in which sums of two squares have square roots). Then we have a constructive version of Tarski's representation theorem:

**THEOREM 9.8.** *(i) The models of intuitionistic Tarski geometry with two-point line-circle continuity are the planes  $F^2$  over a Euclidean field  $F$ .*

*(ii) The models of intuitionistic Tarski geometry with no continuity axioms are the planes  $F^2$  over a Pythagorean field  $F$ .*

*(iii) Given a model of geometry, the field  $F$  and its operations can be explicitly and constructively defined.*

*Proof (of both theorems).* Once we have (uniform) perpendiculars and midpoints, the classical constructions of Descartes and Hilbert that define addition and multiplication can be carried out. In this paper we have proved the existence of (uniform) perpendiculars and midpoints in intuitionistic Tarski geometry, and the definitions of (signed uniform) addition and (signed uniform) multiplication are given in [5]. The field laws are proved for these definitions in [19]; since these are quantifier free when expressed with Skolem functions, they are provable in intuitionistic Tarski geometry (without any continuity axiom) by the double-negation interpretation. The simple construction of the uniform perpendicular suffices for (i), with two-point line-circle, but for (ii) we need the more complicated construction given above, based on Gupta's perpendiculars. Moreover, even for Tarski's (i), with segment-circle in place of two-point line-circle, we need Gupta. That completes the proof.

With classical logic, the representation theorem gives a complete characterization of the consequences of the axioms, because according to Gödel's completeness theorem, a sentence of geometry true in all models  $F^2$  is provable in the corresponding geometry. With intuitionistic logic, the completeness theorem is not valid. The "correct" way to obtain a complete characterization of the theorems of constructive geometry in terms of field theory is to exhibit explicit interpretations mapping formulas of geometry to formulae of field theory, and an "inverse" interpretation from field theory to geometry. We have actually carried this program out, but it is highly technical due to the differences in the two languages, and requires many pages, so we omit it here.

**9.6. Historical Notes.** Tarski claimed in 1959 [24] (page 22, line 5) that he could define addition and multiplication geometrically, and prove the field laws, without using his continuity axiom; hence all models of the axioms (excluding continuity) are planes over ordered fields. The first published proof of these claims was in 1983 [19], and relies heavily on Gupta's 1965 dissertation. In this note we consider what Tarski may have had in mind in 1959.

To define multiplication, we need perpendiculars and midpoints, which were constructed in (A1)-(A10) by Gupta in 1965. Tarski lectured in 1956 on geometry, but I have not found a copy of his lecture notes. He lectured again on geometry in 1963, but according to Gupta did not base his lectures on his own axiom system (and again there are no surviving notes). The proofs in this

section show that he could well have defined addition and multiplication using only two-point line-circle continuity, since we showed here how to construct perpendiculars and midpoints from two-point line-circle continuity. But he made two claims that we do not see how to prove without Gupta: that he could use segment-circle continuity in (i), and that he could get by without any continuity in (ii).

These claims could not have been valid in 1959, since this was some years before Gupta's proofs. In 1959, there was no way known to construct dropped perpendiculars using (A1)-(A10), even with the use of segment-circle continuity; Tarski could not have justified Euclid I.12 on the basis of segment-circle continuity, since it requires the triangle inequality, which requires perpendiculars, and Tarski was not using the triangle circumscription axiom but his own form of the parallel axiom, so he could not even prove that every segment has a midpoint, as far as I can see.

Tarski did believe that line-circle continuity could be derived from a single instance of the continuity schema. He explicitly claimed this in [24], page 26, line 8. But the "obvious" derivation requires the triangle inequality, which in turn seems to require perpendiculars.<sup>20</sup> After Gupta, perpendiculars exist, and the proof of the triangle inequality follows easily, so indeed line-circle continuity follows from A11. But in 1959, there was no known way to get perpendiculars, and so, no way to derive the triangle inequality, and hence, no way to derive line-circle continuity from the continuity schema, or to justify Euclid I.12 to get dropped perpendiculars from line-circle continuity. All these difficulties disappeared once Gupta proved the existence of dropped perpendiculars in (A1)-(A8). Thus Gupta's work has a much more central place than is made apparent in [19]. Tarski desperately needed perpendiculars.

It is possible that Tarski had in mind using two-point line-circle continuity to justify dropped perpendiculars, and overlooked the difficulty of proving two-point line-circle from segment-circle. Even so, to complete a proof of his representation theorem about Euclidean fields, he would have had to duplicate the results in this paper about getting erected perpendiculars and midpoints from dropped perpendiculars. Part (ii) of his theorem, about what happens with no continuity at all, flat-out requires Gupta's work, which put everything right, substantiating the claims of 1959. A discovery of Tarski's missing 1956 lecture notes seem to be the only way to resolve the question of "what Tarski knew and when he knew it."

**§10. Geometry with terms for the intersections of lines.** It seems more natural, when thinking of straightedge-and-compass constructions, to include a symbol  $il(a, b, c, d)$  for the (unique) intersection point of *Line*( $a, b$ ) and

---

<sup>20</sup>It is straightforward to derive line-circle continuity from A11 together with the facts that the interior and exterior of a circle are open, and the interior is convex. These facts in turn are easy to derive from inner Pasch, Euclid III.2 (chord lies inside circle), the density lemma (Lemma 2.7), and the triangle inequality.

*Line*( $c, d$ ). We say “unique” because we want the intersection point of two coincident lines to be undefined; otherwise  $il(a, b, c, d)$  will not be continuous on its domain.

The difficulty with using this Skolem symbol is that the definedness condition for  $il(a, b, c, d)$  is not easily expressible in quantifier-free form. Of course we need  $a \neq b \wedge c \neq d$ , and we want  $\neg(Col(a, b, c) \wedge Col(a, b, d))$  as just explained. But in addition there are parallel lines that do not meet. Using the strong parallel postulate, one can indeed express the definedness condition for  $il(a, b, c, d)$  in a quantifier-free way, namely,  $il(a, b, c, d)$  is defined if and only if there is a point  $p$  collinear with  $a$  and  $b$  but not  $c$  and  $d$ , and a point  $q$  collinear with  $c$  and  $d$ , such that the transversal  $pq$  of the *Line*( $a, b$ ) and *Line*( $c, d$ ) makes alternate interior angles unequal. This condition can be expressed using points only, as shown in Fig. 4 above. We can use the strong parallel axiom to prove stability:

$$\neg\neg il(a, b, c, d) \downarrow \rightarrow il(a, b, c, d) \downarrow .$$

But one cannot do this for subtheories with no parallel postulate or other versions of the parallel postulate. Therefore we prefer, when working with  $il(a, b, c, d)$  to use the Logic of Partial Terms (described below), in which  $t \downarrow$  is made into an official atomic formula for each term  $t$ , instead of an abbreviation at the meta-level.

**10.1. Logic of Partial Terms (LPT).** This is a modification of first-order logic, in which the formation rules for formulas are extended by adding the following rule: If  $t$  is a term then  $t \downarrow$  is a formula. Then in addition the quantifier rules are modified so instead of  $\forall x A(x) \rightarrow A(t)$  we have  $\forall x (t \downarrow \wedge A(x)) \rightarrow A(t)$ , and instead of  $A(t) \rightarrow \exists x A(x)$  we have  $A(t) \wedge t \downarrow \rightarrow \exists x A(x)$ . Details of **LPT** can be found in [2], p. 97.

**LPT** includes axioms  $c \downarrow$  for all constants  $c$  of any theory formulated in **LPT**; this is in accordance with the philosophy that terms denote things, and while terms may fail to denote (as in “the King of France”), there is no such thing as a non-existent thing. Thus  $1/0$  can be undefined, i.e. fail to denote, but if a constant  $\infty$  is used in **LPT**, it must denote something.

The meaning of  $t = s$  is that  $t$  and  $s$  are both defined and they are equal. We write  $t \cong s$  to express that if one of  $t$  or  $s$  is defined, then so is the other, and they are equal.

**DEFINITION 10.1.** *For terms in any theory using the logic of partial terms,  $t \cong q$  means*

$$t \downarrow \rightarrow t = q \wedge q \downarrow \rightarrow t = q.$$

*This is read  **$t$  and  $q$  are equal if defined.***

Thus “ $\cong$ ” is an abbreviation at the meta-level, rather than a symbol of the language.

**LPT** contains the axioms of “strictness”, which are as follows (for each function symbol  $f$  and relation symbol  $R$  in the language):

$$\begin{aligned} f(t_1, \dots, t_n) \downarrow &\rightarrow t_1 \downarrow \wedge \dots \wedge t_n \downarrow \\ R(t_1, \dots, t_n) &\rightarrow t_1 \downarrow \wedge \dots \wedge t_n \downarrow \end{aligned}$$

Note that in **LPT**, under a given “valuation” (assignment of elements of a structure to variables), each formula has a definite truth value, i.e. we do not use three-valued logic in the semantics. For example, if  $P$  is a formula of field theory with a reciprocal operation  $1/x$ , then  $P(1/0)$  is false, since  $1/0$  is undefined. For the same reason  $\neg P(1/0)$  is false. Hence  $P(1/0) \vee \neg P(1/0)$  is false too; but that does not contradict the classical validity of  $\forall x(P(x) \vee \neg P(x))$  since we are required to prove  $t \downarrow$  before deducing an instance  $P(t) \vee \neg P(t)$ .

As an example of the use of **LPT**, we reformulate the theory of Euclidean fields [5] using the logic of partial terms. The existential quantifiers associated with the reciprocal axioms, with the axiom of additive inverse, and with the square-root axiom of Euclidean field theory are replaced by a function symbol  $\sqrt{\phantom{x}}$ , a unary minus  $-$ , and a function symbol for “reciprocal”, which we write as  $1/x$  instead of *reciprocal*( $x$ ). The changed axioms are

$$\begin{aligned} x + (-x) &= 0 && \text{(additive inverse)} \\ x \neq 0 &\rightarrow x \cdot (1/x) = 1 && \text{(EF1')} \\ P(x) &\rightarrow x \cdot (1/x) = 1 && \text{(EF7')} \\ x + y = 0 \wedge \neg P(y) &\rightarrow \sqrt{x} \cdot \sqrt{x} = x && \text{(EF5')} \end{aligned}$$

**10.2. A version of Tarski’s theory with terms for intersections of lines.** This version of Tarski’s theory we call ruler-and-compass Tarski. It is formulated as follows:

- It uses a function symbol  $il(a, b, p, q)$  for the intersection point of  $Line(a, b)$  and  $Line(p, q)$ .
- It uses the logic of partial terms.
- If  $il(a, b, p, q)$  is defined, then it is a point on both lines.
- If there is a point on  $Line(a, b)$  and  $Line(p, q)$ , and those lines do not coincide, then  $il(a, b, p, q)$  is such a point.
- Formally, the axioms involving  $il$  are

$$Col(a, b, x) \wedge Col(p, q, x) \wedge \neg(Col(a, b, p) \wedge Col(a, b, q)) \rightarrow il(a, b, p, q) \downarrow$$

$$il(a, b, p, q) \downarrow \rightarrow a \neq b \wedge p \neq q \wedge Col(a, b, il(a, b, p, q)) \wedge Col(p, q, il(a, b, p, q))$$

- $il$  is used instead of separate Skolem functions for  $ip$ . Specifically, the term  $ip(a, p, c, b, q)$  in the Skolemized inner Pasch axiom become  $il(a, q, b, p)$ . The point  $c$  does not occur in this term.
- The Skolem functions *ext* (for segment extension) is not changed.
- The Skolem functions for intersections of lines and circles are not changed.
- Stability for equality, betweenness, and congruence, as before.

We could consider replacing Skolem terms  $center(a, b, c)$  with terms built up from  $il$ . The two lines to be intersected are the perpendicular bisectors of  $ab$  and  $bc$ , where  $a, b$ , and  $c$  are three non-collinear points. We can define the perpendicular bisector by the erected perpendicular at the midpoint, so it is indeed possible to eliminate the symbol *center*; but there seems to be no special reason to do so.

We did not include the stability of definedness; that is because it can be proved. The following lemma is proved in [5]; here we give a different proof, based on the triangle-circumscription form of the strong parallel axiom.

LEMMA 10.2. [Stability of  $il(a, b, c, d)$ ] *The strong parallel postulate is equivalent (in ruler-and-compass Tarski minus the parallel postulate) to the stability of  $il(a, b, c, d) \downarrow$ :*

$$\neg\neg il(a, b, c, d) \downarrow \rightarrow il(a, b, c, d) \downarrow .$$

*Proof.* (i) First suppose the strong parallel postulate and  $\neg\neg il(a, b, c, d) \downarrow$ . We will show  $il(a, b, c, d) \downarrow$ . Let  $L = Line(a, b)$  and  $K = Line(c, d)$ . Then lines  $K$  and  $L$  do not coincide, for then  $il(a, b, c, d)$  would be undefined. Hence we can find a point on  $L$  that is not on  $K$ . We may assume without loss of generality that  $b$  is such a point. Construct point  $f$  so that  $bf$  is parallel to  $K$ ; more explicitly,  $K$  and  $bf$  and the transversal  $bc$  make alternate interior angles equal. If  $a, b$ , and  $f$  are collinear, then  $ab$  and  $cd$  are parallel, so  $il(a, b, c, d)$  is undefined, contradiction. Hence  $a, b$ , and  $f$  are not collinear. Then line  $M = Line(b, f)$  passes through point  $b$  and is parallel to  $K$ , and line  $L$  also passes through  $b$ , and has a point  $a$  not on  $M$ . Then by the strong parallel axiom,  $L$  meets  $K$ . In that case  $il(a, b, c, d)$  is defined, as claimed.

(ii) Conversely, suppose the stability of  $il(a, b, c, d) \downarrow$ , and suppose  $a, b$ , and  $c$  are not collinear. Let  $m$  be the midpoint of  $ab$  and  $n$  the midpoint of  $cd$ , with  $pm$  the perpendicular bisector of  $ab$  and  $qn$  the perpendicular bisector of  $cd$ . We must prove  $il(m, p, n, q) \downarrow$ . By stability it suffices to derive a contradiction from the assumption that it is not defined. If it is not defined then  $mp$  is parallel to  $nq$  (as not meeting is the definition of parallel). But  $Line(a, b)$  and  $Line(b, c)$  are perpendicular to  $mp$  and  $nq$  respectively; hence they cannot fail to be parallel or coincident. But since they both contain point  $b$ , they are not parallel; hence they are coincident. Hence  $a, b$ , and  $c$  are collinear, contradiction. That completes the proof.

THEOREM 10.3 (Stability of definedness). *For each term  $t$  of ruler-and-compass Tarski,  $\neg\neg t \downarrow \rightarrow t \downarrow$  is provable.*

*Proof.* By induction on the complexity of the term  $t$ . If  $t$  is a compound term  $ts$ , and  $\neg\neg ts \downarrow$ , then  $\neg\neg t \downarrow$  and  $\neg\neg s \downarrow$ , so by induction hypothesis,  $t \downarrow$  and  $s \downarrow$ . Hence  $ts \downarrow$ . We may therefore suppose  $t$  is not a compound term. If the functor is  $ic_1, ic_2, ilc_1$ , or  $ilc_2$ , then it is easy to prove that the conditions for  $t$  to be defined are given geometrically, by the same formulas that were used to define  $t \downarrow$  in Tarski with Skolem functions (and without **LPT**). Hence stability follows by the stability of equality, congruence, and betweenness. The stability of  $il(a, b, c, d)$  is equivalent to the strong parallel postulate, by the previous lemma. That completes the proof.

**10.3. Intersections of lines and the parallel axiom.** In the proof of the first part of Lemma 10.2, we showed that if lines  $L$  and  $M$  meet in a point  $x$ , then  $x$  can be made to appear as the center of a circle circumscribed about suitably chosen points  $a, b$ , and  $c$ . In this section, we will refine this construction to show that there is a single term  $t(a, b, c, d)$  in the language of Tarski with Skolem functions that give the intersection point of  $Line(a, b)$  and  $Line(c, d)$ , when it exists.

LEMMA 10.4. *Given two lines  $L$  and  $K$  that are neither coincident nor parallel, one can construct a point  $p$  that lies on  $K$  but not on  $L$ . More precisely,*

interpreting  $L$  as  $Line(a, b)$  and  $K$  as  $Line(q, r)$ , there is a single term  $t(a, b, q, r)$  such that if  $\neg(Col(a, b, q) \wedge Col(a, b, r))$  and for some  $x$ ,  $Col(a, b, x) \wedge Col(q, r, x)$ , then  $e = t(a, b, q, r)$  satisfies  $Col(q, r, e) \wedge \neg Col(a, b, e)$ .

*Remarks.* The point  $x$  cannot be used to construct  $e$ , which must depend only on  $a, b, q$ , and  $r$ , and must be constructed by a single term, and hence depend continuously on the four parameters. We will use the parallel postulate to construct  $e$ ; we do not know a construction that does not use the parallel postulate.

*Proof.* Let  $M$  be the perpendicular to  $K$  passing through  $q$ . We are supposed to construct  $M$  from  $a, b, q$ , and  $r$  alone. To construct  $M$ , we need not just  $p$  and  $q$ , but also a point not on  $K$ ; and  $a$  and  $b$  are useless here as they might lie on  $K$ . We must appeal to Lemma 11.7 for the construction of some point not on line  $K$ ; thus this apparently innocent lemma requires the geometric definition of arithmetic and the introduction of coordinates, and hence the parallel postulate.

The construction of  $M$  gives us one point  $u$  on  $M$  different from  $q$ . Let  $v$  be the reflection of  $u$  in  $q$ . Then  $u$  and  $v$  are equidistant from  $q$ . Now, using the uniform perpendicular construction we construct the line  $J$  through  $u$  perpendicular to  $L$ . See Fig. 23.<sup>21</sup>

While we do not know whether  $u$  lies on  $L$ , the uniform perpendicular construction (Theorem 9.5) provides two points determining  $J$ , namely

$f = Project(u, a, b, c)$  and  $h = head(u, a, b, c)$ , where  $f$  is on  $L$  and  $c$  is not on  $L$ . Possibly  $u$  is equal to  $f$  or to  $h$ ; we need a point  $d$  on  $J$  that is definitely not equal to  $u$  or to the reflection of  $u$  in  $L$ . To get one, our plan is to draw a circle of sufficiently large radius about  $u$  and intersect it with  $J = Line(f, h)$ . We use the uniform reflection construction to define  $w = Reflect(u, a, b)$ , the reflection of  $w$  in  $L$ . Then we extend the non-null segment  $\alpha\beta$  by the (possibly null) segment  $uw$  to get a point  $z$  such that  $\alpha z > uw$ . Then we use  $\alpha z$  as the "sufficiently large" radius. Here is the construction:

$$\begin{aligned} z &= ext(\alpha, \beta, u, w) \\ d &= ilc_1(f, c, u, e_2(u, \alpha, z)) \end{aligned}$$


Now  $d$  lies on  $J$  and is different from  $u$ , and it is also different from  $w$  since  $w$  lies inside the circle centered at  $u$  of radius  $\alpha z$ . Finally define

$$c = center(u, v, d).$$

The three points  $u, v$ , and  $d$  are not collinear, since then  $J$  and  $M$  would coincide, and  $L$  and  $K$  would both be perpendicular to  $J$ , and hence parallel; but  $L$  and  $K$  are by hypothesis not parallel. Since  $u, v$ , and  $d$  are not collinear,  $c$  is equidistant from  $u, v$ , and  $d$ . Therefore  $c$  lies on the perpendicular bisector of  $uv$ , which is  $K$ . Also  $c$  lies on the perpendicular bisector of  $ud$ , which is parallel to  $L$ , since both are perpendicular to  $J$ . This perpendicular bisector does not coincide with

<sup>21</sup>In a theory with Skolem functions for the intersection points of two circles, the construction of  $M$  and  $u$  becomes trivial (just use the Euclidean construction of a perpendicular), but axiomatizing the Skolem functions for the circle in such a way as to distinguish the points of intersection (which is necessary to construct perpendiculars) requires the introduction of coordinates. So using circle-circle does not obviate the need for coordinates in this lemma.

FIGURE 23. Uniform construction of a point  $c = \text{center}(d, u, v)$  on  $K$  that is not on  $L$ . The construction works whether or not  $q$  is on  $L$ , or  $u$  is on  $L$ . The dotted line bisects  $ud$  and does not coincide with  $L$ .



$L$ , since  $d$  is not the reflection  $w$  of  $u$  in  $L$ . Therefore  $c$  does not lie on  $L$ . Then  $c$  lies on  $K$  but not on  $L$ , as desired. That completes the proof of the lemma.

**THEOREM 10.5.** [*Elimination of  $il$* ] There is a term  $t(a, b, p, r)$  of intuitionistic Tarski with Skolem functions (so  $t$  contains  $ip$  and  $\text{center}$  but not  $il$ ) such that the following is provable:

$$\text{Col}(a, b, x) \wedge \text{Col}(p, r, x) \wedge \neg(\text{Col}(a, b, p) \wedge \text{Col}(a, b, r)) \wedge p \neq r \rightarrow x = t(a, b, p, r).$$


In other words,  $t(a, b, p, r)$  gives the intersection point of  $\text{Line}(a, b)$  and  $\text{Line}(p, r)$ .

*Remark.* The problem here is to explicitly produce the term  $t$  that is implicit in the proof of Lemma 10.2. This is also closely related to the proof given in [5] that in constructive neutral geometry, the triangle circumscription principle implies the strong parallel axiom. But here we have to check that this construction can be carried out in Tarski's geometry, i.e., all the lines that are required to intersect are proved to intersect using only inner Pasch; so there is definitely something additional to check.

*Proof.* By Lemma 10.4, we may assume without loss of generality that  $p$  does not lie on  $L$ . More explicitly, if we now produce a term  $t$  as in the lemma, that works under the additional assumption that  $p$  is not on  $L$ , then we can compose that term with the term given in Lemma 10.4, which produces a point on  $\text{Line}(p, r)$  that is not on  $L$ , and the composed term will work without the assumption that  $p$  is not on  $L$ .

Recall that  $Project(p, a, b)$  is the point  $w$  on  $Line(a, b)$  such that  $pw \perp ab$ , and  $Project(p, a, b)$  is given by a term in Tarski with Skolem functions. There is also a term  $erect(p, r)$  that produces a point  $j$  such that  $j$  is not on  $Line(p, r)$  and  $jp \perp pr$ . (Since  $p$  is on  $Line(p, r)$ , the uniform perpendicular construction is not needed; the simple Euclidean construction will be enough.) Finally, there is a term  $Reflect(x, a, b)$  that produces the reflection of  $x$  in  $Line(a, b)$ ; if we assume  $x$  is not on  $Line(a, b)$  this is particularly easy: let  $q = Project(x, a, b)$  and take  $Reflect(x, a, b) = ext(x, q, x, q)$ . The requirement that  $x$  is not on  $Line(a, b)$  means that we are extending a non-trivial segment, which is constructively allowed.

FIGURE 24. Triangle circumscription implies the strong parallel axiom. Given lines  $L$  and  $M$ , to construct their intersection point as the center  $e$  of an appropriate circle.  $y$  and  $z$  are reflections of  $x$  in  $M$  and  $L$ .



In Fig. 24,  $L = Line(a, b)$  and  $M = Line(p, r)$ , and  $p$  does not lie on  $L$ . First we claim that  $x$  does not lie on  $L$ . If  $x$  does lie on  $L$ , then  $px$  is a right triangle, so the hypotenuse  $px$  is greater than the leg  $pw$ ; but  $px = mx$ , which is less than  $mw$  since  $m = midpoint(p, w)$ , and  $p \neq w$ . Therefore  $Reflect(x, a, b)$  can be defined using the easy construction given above.

Now we give a construction script corresponding to the figure:

$$\begin{aligned}
 w &= Project(p, a, b) \\
 m &= midpoint(p, w) \\
 j &= erect(p, r) \\
 x &= ilc_1(j, p, p, m) \\
 y &= ilc_2(j, p, p, m) \\
 z &= Reflect(x, a, b) \\
 e &= center(x, y, z)
 \end{aligned}$$

Composing the terms listed above we find a (rather long) term that produces  $e$  from  $a$ ,  $b$ ,  $p$ , and  $r$ . We claim that  $e$  is the intersection point of  $Line(a, b)$  and  $Line(p, r)$ . By the stability of collinearity, we can argue by cases on whether  $x$ ,  $y$ , and  $z$  are collinear or not. (Here it is important that we are not proving a statement with an existential quantifier, but a quantifier-free statement involving a single term that constructs the desired point  $e$ .) In case  $x$ ,  $y$ , and  $z$  are not collinear, then  $center(x, y, z)$  produces a point  $e$  that is the center of a circle containing  $x$ ,  $y$ , and  $z$ . Then Euclid III.1 implies that  $L$  and  $M$  both pass through  $e$ , and we are done. On the other hand, if  $x$ ,  $y$ , and  $z$  are collinear, then  $M$  and  $L$  are both perpendicular to  $Line(x, y)$ , so  $M$  and  $L$  are parallel; then there is no  $x$  as in the hypothesis of the formula that is alleged, so there is nothing more to prove. That completes the proof of the lemma.

**10.4. Interpreting ruler-and-compass Tarski in intuitionistic Tarski with Skolem functions.** Ruler-and-compass Tarski clearly suffices to interpret intuitionistic Tarski (with or without Skolem functions), because the points asserted to exist by inner Pasch and the triangle circumscription principle are given as intersections of lines. Conversely we may ask, whether ruler-and-compass Tarski can be interpreted in intuitionistic Tarski with Skolem functions. That is, can terms in  $il$  be effectively replaced by terms built up from  $ip$  and  $center$ ? The answer is “yes”.

**THEOREM 10.6.** *Suppose ruler-and-compass Tarski geometry (with  $il$  and other Skolem functions) proves a theorem  $\phi$  that does not contain  $il$ . Then Tarski geometry with Skolem functions proves  $\phi$ . Moreover if  $\phi$  contains no Skolem functions, then Tarski geometry proves  $\phi$ . These claims hold both for the theories with intuitionistic logic and those with classical logic.*

*Proof.* We assign to each term  $t$  of ruler-and-compass Tarski, a corresponding term  $t^\circ$  of intuitionistic Tarski with Skolem functions. Let  $il^\circ(a, b, q, r)$  be the term given by Theorem 10.5. The term  $t^\circ$  is defined inductively by

$$\begin{aligned} x^\circ &= x && \text{where } x \text{ is a variable or constant} \\ il^\circ(a, b, c, d) &= il^\circ(a^\circ, b^\circ, c^\circ, d^\circ) \\ f(a, b, c, d) &= f(a^\circ, b^\circ, c^\circ, d^\circ) && \text{where } f \text{ is } ilc_1, ilc_2, ic_1, ic_2, \text{ or } ext \\ e(a)^\circ &= e(a^\circ) \end{aligned}$$

Then we assign to each formula  $A$  of ruler-and-compass Tarski, a corresponding formula  $A^\circ$  of intuitionistic Tarski with Skolem functions. Namely, the map  $A \mapsto A^\circ$  commutes with logical operations and quantifiers, and for atomic  $A$  not of the form  $t \downarrow$ , we define

$$A(t_1, \dots, t_n)^\circ = A(t_1^\circ, \dots, t_n^\circ).$$

For the case when  $A$  is  $t \downarrow$ , we define  $(t \downarrow)^\circ$  to be  $t = t$  when  $t$  is a variable or constant, and when it is a compound term, we use Definition 5.2. By induction on the complexity of  $A$ , we see that  $A^\circ[x := t^\circ]$  is provably equivalent to  $(A[x := t])^\circ$ .

Then by induction on the length of proofs in ruler-and-compass Tarski, we show that if ruler-and-compass Tarski proves  $\phi$ , then intuitionistic Tarski with Skolem functions proves  $\phi^\circ$ . A propositional axiom or inference remains one

under the interpretation, so it is not even vital to specify exactly which propositional axioms we are using. In this direction (from **LPT** to ordinary logic), the quantifier rules and axioms need no verification, as the extra conditions of definedness needed in **LPT** are superfluous in ordinary logic. For example, one of those axioms is  $\forall x A \wedge t \downarrow \rightarrow A[x := t]$ . That becomes  $\forall x A^\circ \wedge (t \downarrow)^\circ \rightarrow A[x := t^\circ]$ , in which the  $t \downarrow$  can just be dropped. There are some special axioms in **LPT**, for example  $c \downarrow$  for  $c$  a constant and  $x \downarrow$  for  $x$  a variable.

We check the basic axioms for  $ilc_1$ . These say that (i) if  $ilc_1(a, b, c, d)$  is defined, then it is a point on  $Line(a, b)$  and also on the circle with center  $c$  passing through  $d$ , and (ii) if there is a point  $x$  on both the line and circle, then  $ilc_1(a, b, c, d)$  is defined. According to the definition of  $(ilc_1(a, b, c, d) \downarrow)^\circ$ , the interpretation of “ $ilc_1 \downarrow$ ” is “there is a point on the line inside the circle”, where “inside” means not strictly inside. Since  $ilc_1$  is not affected by the interpretation (except in the atomic formula  $ilc_1(a, b, c, d) \downarrow$ ), the interpretations of the basic axioms for  $ilc_1$  are equivalent to those same axioms. Similarly for  $ilc_2$ ,  $ic_1$ , and  $ic_2$ .

Now consider the axioms for  $il(a, b, p, r)$ . These axioms say that if  $il(a, b, p, r) \downarrow$  then  $e = il(a, b, p, r)$  is a point on  $Line(a, b)$  and also on  $Line(p, r)$ , and if  $x$  is any point on both lines and not both  $p$  and  $r$  lie on  $Line(a, b)$ , then  $x = il(a, b, p, r)$ . Recall that  $(il(a, b, p, r) \downarrow)^\circ$  says there exists an  $x$  on both lines, and not both  $p$  and  $r$  are on  $Line(a, b)$ . Then the interpretation of these axioms says that if  $Line(a, b)$  and  $Line(p, r)$  meet and not both  $p$  and  $r$  lie on  $Line(a, b)$ , then  $il(a, b, p, r)^0$  is the intersection point. But that is exactly Theorem 10.5. That completes the proof.

**THEOREM 10.7.** *Suppose classical ruler-and-compass Tarski geometry proves a theorem of the form  $P \rightarrow Q(t)$ , with  $P$  and  $Q$  negative. Then intuitionistic Tarski geometry proves  $P \rightarrow \exists x Q(x)$ .*

*Proof.* Suppose  $P \rightarrow Q(t)$  is provable in intuitionistic ruler-and-compass geometry. Then  $P \rightarrow \exists x Q(x)$  is provable in intuitionistic ruler-and-compass geometry. But that formula contains no occurrences of  $il$ . Then by Theorem 10.6, it is provable in intuitionistic Tarski geometry. That completes the proof.

**COROLLARY 10.8** (Pasch's theorem). *The following version of Pasch's axiom is provable in intuitionistic Tarski: given triangle  $abc$ , and line  $L$  not containing  $a$ ,  $b$ , or  $c$ , and suppose that  $L$  meets  $ab$  but not  $bc$ . Then  $L$  meets  $ac$ .*

*Proof.* Let  $L = Line(u, v)$ , and let  $p$  lie on  $L$  with  $\mathbf{B}(a, p, b)$ . Since Pasch's axiom is provable in classical Tarski geometry, it is also provable in classical ruler-and-compass Tarski geometry. It has the form  $P \rightarrow \exists x Q(x)$ , where  $P$  says that  $Line(u, v)$  does not contain  $a$ ,  $b$ , or  $c$ , and meets  $ab$  but not  $bc$ , and  $Q(x)$  is  $\mathbf{B}(a, x, c) \wedge Col(u, v, x)$ . But in ruler-and-compass geometry, the point  $x$  asserted to exist is given as the intersection point of  $L$  and  $Line(a, c)$ , namely  $il(u, v, a, c)$ . Then Theorem 10.7 tells us that intuitionistic Tarski geometry proves  $P(x) \rightarrow \exists x Q(x)$ , which is the desired version of Pasch's axiom. That completes the proof of the corollary.

*Remark.* This result at first seems rather remarkable, as the classical proof of Pasch from inner Pasch and outer Pasch uses an argument by cases. But we can

sketch a direct constructive proof of Pasch's theorem as follows. Note the role played by the strong parallel axiom in the proof.

There are three cases:  $L$  parallel to  $bc$ , or  $L$  meets  $bc$  at  $x$  with  $\mathbf{B}(b, c, x)$ , or  $L$  meets  $bc$  at  $x$  with  $\mathbf{B}(x, b, c)$ . If  $L$  does not meet  $ac$ , then the second and third cases contradict inner Pasch and outer Pasch, respectively, and the first case results in two parallels to  $L$  through  $c$ , which is contradictory. Hence  $L$  cannot fail to meet  $ab$ . Let  $L = \text{Line}(u, v)$ ; so  $\neg\neg il(u, v, a, c) \downarrow$ . By the stability of  $il(u, v, a, c) \downarrow$ , we have  $il(u, v, a, c) \downarrow$ . Let  $x = il(u, v, a, c)$ . Then we only have to show  $\mathbf{B}(a, x, c)$ . But if  $\neg\mathbf{B}(a, x, c)$ , then similarly  $L$  cannot fail to meet  $bc$ , contradicting the hypothesis. Hence  $\neg\neg\mathbf{B}(a, x, c)$ . Hence by the stability of betweenness,  $\mathbf{B}(a, x, c)$ .

*Remark.* Theorem 10.7 settles the question of whether the version of Pasch given in (A7-i) is sufficiently general. We could, for example, have considered a more general version allowing for point  $p$  to be on  $\text{Ray}(a, c)$  and not equal to  $a$ ; outer Pasch covers the case when  $\mathbf{T}(a, c, p)$  and inner Pasch covers the case when  $\mathbf{T}(a, p, c)$ , but constructively these cases are not exhaustive. (What about a point  $p$  whose order relations to  $c$  are unknown?) But the corresponding classical existence theorem is provable (by cases), so by the theorem, it is already provable in intuitionistic Tarski geometry.

In fact, the theorem shows that we could just as well have restricted (A7-i) further by using  $\mathbf{B}$  instead of  $\mathbf{T}$ , as that would still suffice to construct perpendiculars and midpoints, which are all we need to be able to use *center* to find intersection points of lines.

*Remark.* If we drop the strong parallel axiom (or triangle superscription principle), we obtain "neutral geometry with  $il$ ". It is an open question whether neutral geometry with  $il$  can be interpreted in neutral Tarski with Skolem functions. In other words, can all terms for intersection points of lines that are needed in proofs of theorems not mentioning  $il$  be replaced by terms built up from  $ip$  and  $ext$ ? We used *center* in an essential way in the proof of Theorem 10.5, but did we have to do so? We think there is probably no way to do this without using the parallel postulate.

**§11. Relations between classical and constructive geometry.** Our intuition about constructive geometry is this: You may argue classically for the equality or inequality of points, for the betweenness of points, for collinearity, for the congruence of segments. But if you assert that something exists, it must be constructed by a single, uniform construction, not by different constructions applying in different cases. If you can give a uniform construction, you may argue by cases that it works, but the construction itself cannot make a case distinction. Thus the uniform perpendicular construction of a line through  $x$  perpendicular to  $L$  works whether or not  $x$  is on  $L$ ; if we wished, we could argue for its *correctness* by cases, as we could always push a double negation through the entire argument and use stability to eliminate it.

There is in Szmielew's Part I of [19] an extensive development from Tarski's classical axioms, essentially deriving Hilbert's axioms and the definitions and

key properties of addition and multiplication. We would like to be able to import arguments and results wholesale from this development into constructive geometry. In this section we investigate to what extent this is possible.

It is certainly not completely possible to import results without modifying Tarski's axioms, since constructive proofs will produce points that depend continuously on parameters, while as we have discussed above, Tarski's version of inner Pasch and segment extension axioms do not have this property. Those defects have been remedied above by formulating "continuous Tarski geometry", a theory classically equivalent to Tarski's geometry.

**11.1. The double-negation interpretation.** Gödel introduced [9] his double-negation interpretation, which assigns a formula  $A^-$  to every formula  $A$ , by replacing  $\exists$  by  $\neg\forall\neg$  and replacing  $A \vee B$  by  $\neg(\neg A \wedge \neg B)$ . For atomic formulae,  $A^-$  is defined to be  $\neg\neg A$ . The rules of intuitionistic logic are such that if  $A$  is classically provable (in predicate logic) then  $A^-$  is intuitionistically provable. Hence, if we have a theory  $T$  with classical logic, and another theory  $S$  with intuitionistic logic, whose language includes that of  $T$ , and for every axiom  $A$  of  $T$ ,  $S$  proves  $A^-$ , then  $S$  also proves  $A^-$  for every theorem  $A$  of  $T$ . In case the atomic formulas in the language of  $T$  are stable in  $S$ , i.e. equivalent to their double negations, then of course we can drop the double negations on atomic formulas in  $A^-$ .

In [3] we applied this theorem to a version of constructive geometry based on Hilbert's axioms. Given the extensive almost-formal development of geometry from Tarski's axioms in [19], one might like to use the double-negation interpretation with  $T$  taken to be Tarski's theory, and  $S$  taken to be some suitable constructive version of Tarski's theory. We now investigate this possibility.

A double-negation interpretation from a classical theory to a constructive version of that theory becomes a better theorem if it applies to the Skolemized versions of the theories, because in the un-Skolemized version, an existential quantifier is double-negated, while the corresponding formula of the Skolemized theory may replace the existentially quantified variable by a term, so no double negated quantifier is involved, and no constructive content is lost. But if we Skolemize Tarski's version of inner Pasch, we get an essentially non-constructive axiom, as shown above. Hence there is no double-negation interpretation for that theory. However, it works fine if we replace Tarski's axioms by the (classically equivalent) axioms of continuous Tarski with Skolem functions:

**THEOREM 11.1.** *Let  $T$  be intuitionistic Tarski geometry with Skolem functions. If  $T$  plus classical logic proves  $\phi$ , then  $T$  proves the double-negation interpretation  $\phi^-$ .*

*Proof.* It suffices to verify that the double-negation interpretations of the axioms are provable. But the axioms are negative and quantifier-free, so they are their own double-negation interpretations. That completes the proof.

**COROLLARY 11.2.** *If  $\phi$  is negative, and classical Tarski geometry without Skolem functions proves  $\phi$ , then intuitionistic Tarski geometry proves  $\phi$ .*

*Proof.* Suppose  $\phi$  is provable in classical Tarski geometry (with or without Skolem functions). Then since  $\phi$  itself has no Skolem functions,  $\phi$  is provable in

classical Tarski geometry without Skolem functions, and hence by Theorem 6.2, it is provable in intuitionistic Tarski geometry with Skolem functions. Hence, by Theorem 11.1,  $\phi^-$  is provable. Since  $\phi$  is negative, it is equivalent to  $\phi^-$ . That completes the proof.

We illustrate the use of Theorem 11.1 by importing the work of Eva Kallin, Scott Taylor, H. N. Gupta, and Tarski mentioned in Section 2.7.

**COROLLARY 11.3.** *The formulas (A16) through (A18), which were once axioms of Tarski's theory, but were shown classically provable from the remaining axioms, are also provable in intuitionistic Tarski without Skolem functions.*

*Proof.* By Corollary 11.2.

We would like to emphasize something has been achieved with the double-negation theorem even for negative theorems, as it would be quite laborious to check the long proofs of (A16)-(A18) directly to verify that they are constructive. For example, (A18) is Satz 5.1 in [19]. Let us consider trying to check directly if this proof is constructive. You can see that the proof proceeds by contradiction, which is permissible by stability; in the crucial part of the proof, inner Pasch is applied to a triangle which ultimately must collapse (as the contradiction is reached) to a single point. Therefore we can constructivize this part provided the non-collinearity hypothesis is satisfied for the application of Pasch. By stability, we may assume that the vertices of this triangle are actually collinear. But can we finish the proof in that case? It looks plausible that (A15) or similar propositions might apply, but it is far from clear. Yet the double-negation interpretation applies, and we do not need to settle the issue by hand. We had to assume (A15), but we do not have to assume (A18), because it is already provable.

The following theorems (numbered as in [19]) have proofs simple enough to check directly (as we did before developing the double negation interpretation), but with the aid of the double negation interpretation, we do not need to check them directly.

**LEMMA 11.4.** *The following basic properties of betweenness are provable in intuitionistic Tarski geometry. Note that  $\mathbf{T}(a, b, c)$  is a defined concept;  $\mathbf{B}(a, b, c)$  is primitive. The theorem numbers refer to [19].*

$\mathbf{T}(a, b, b)$	Satz 3.1
$\mathbf{T}(a, b, c) \rightarrow \mathbf{T}(b, c, a)$	Satz 3.2
$\mathbf{T}(a, a, b)$	Satz 3.3
$\mathbf{T}(a, b, c) \wedge \mathbf{T}(b, a, c) \rightarrow a = b$	Satz 3.4
$\mathbf{T}(a, b, c) \wedge \mathbf{T}(b, c, d) \rightarrow \mathbf{T}(a, b, c)$	Satz 3.5a
$\mathbf{T}(a, b, c) \wedge \mathbf{T}(a, c, d) \rightarrow \mathbf{T}(b, c, d)$	Satz 3.6a
$\mathbf{T}(a, b, c) \wedge \mathbf{T}(b, c, d) \wedge b \neq c \rightarrow \mathbf{T}(a, c, d)$	Satz 3.7a
$\mathbf{T}(a, b, d) \wedge \mathbf{T}(b, c, d) \rightarrow \mathbf{T}(a, c, d)$	Satz 3.5b
$\mathbf{T}(a, b, c) \wedge \mathbf{T}(a, c, d) \rightarrow \mathbf{T}(a, d, b)$	Satz 3.6b
$\mathbf{T}(a, b, c) \wedge \mathbf{T}(b, c, d) \wedge b \neq c \rightarrow \mathbf{T}(a, d, b)$	Satz 3.7b

*Proof.* By the double negation interpretation, since each of these theorems is negative.

Does the double negation interpretation help us to be able to “import” proofs of existential theorems from [19] to intuitionistic Tarski? It gives us the following recipe: Given an existential theorem proved in classical Tarski, we examine the proof to see if we can construct a Skolem term (or terms) for the point(s) asserted to exist. If the proof constructs points using inner Pasch, we need to verify whether degenerate cases or a possibly collinear case are used. If they are not used then the strict inner Pasch axiom (A7-i) suffices. The crucial question is whether the point alleged to exist can be constructed by a single term, or whether the proof is an argument by cases in which different terms are used for different cases. In the latter case, the proof is not constructive (though the theorem might still be, with a different proof). But in the former case, the double-negation interpretation will apply.

Thus the double-negation fully justifies the claim that the essence of constructive geometry is the avoidance of arguments by cases, providing instead uniform constructions depending continuously on parameters.

**11.2. Euclid I.2 revisited.** Consider the first axiom of Tarski’s geometry, which says any segment (null or not) can be extended:  $\exists d(\mathbf{T}(a, b, d) \wedge bd = bc)$ . Clearly  $d$  cannot depend continuously on  $a$  as  $a$  approaches  $b$  while  $b$  and  $c$  remain fixed, since as  $a$  spirals in towards  $b$ ,  $d$  circles around  $b$  outside a fixed circle. Therefore Axiom (A4) of Tarski’s (classical) geometry is essentially non-constructive; the modification to (A4-i) that we made in order to pass to a constructive version was essential.

Euclid I.2 says that given three points  $a, b, c$ , there exists a point  $d$  such that  $ad = bc$ . Euclid gave a clever proof that works when the three points are distinct, and classically a simple argument by cases completes the proof. Constructively, that does not work, since when  $b$  and  $c$  remain fixed and  $a$  approaches  $b$ ,  $d$  from Euclid’s construction does not depend continuously on  $a$ . We will show in this section that it is only Euclid’s proof that is non-constructive; the theorem itself is provable in intuitionistic Tarski geometry, by a different proof.

LEMMA 11.5. *Intuitionistic Tarski geometry proves*

- (i)  $\mathbf{T}(a, b, c)$  and  $\mathbf{T}(p, q, r)$  and  $ac = pr$  and  $bc = qr$  then  $ab = pq$
- (ii) a segment  $ac$  cannot be congruent to (a proper subsegment)  $bc$  with  $\mathbf{B}(a, b, c)$ .

*Proof.* We first show (ii) follows from (i). Suppose  $ac = bc$  and  $\mathbf{B}(a, b, c)$ . Then in (i) take  $p = q = a$  and  $r = c$ . Then (i) implies  $ab = aa$ , contrary to axiom (A3). Hence (i) implies (ii) as claimed.

Now (i) is Satz 4.3 in [19], and since it is negative, we can conclude from the double negation interpretation that it is constructively provable. That completes the proof.

LEMMA 11.6. *In intuitionistic Tarski geometry, null segments can be extended, and Euclid I.2 is provable. Indeed, there is a term (using Skolem functions)  $e(x)$  such that  $e(x) \neq x$  is provable, and a term  $e_2$  corresponding to Euclid I.2, such that if  $d = e_2(a, b, c)$ , then  $ad = bc$ .*

*Remarks.* Thus, it is only Euclid’s proof of I.2 that is non-constructive, as discussed in [5], not the theorem itself. Note that a constructive proof of the theorem

should produce a continuous vector field on the plane, so the constructive content of  $\forall x \exists y (y \neq x)$  is nontrivial. Notice how the proof fulfills this prediction.

*Proof.* Let  $\alpha$  and  $\beta$  be two of the three constants used in the dimension axioms, and define

$$e(x) = ext(\alpha, \beta, \alpha, x)$$

Since  $\alpha \neq \beta$  we have  $e(x) \downarrow$ . Let  $d = e(x)$ ; we claim  $d \neq x$ . By (A4) we have  $\mathbf{T}(\alpha, \beta, d)$  and  $\beta d = \alpha d$ . Then the subsegment  $\beta d$  is congruent to the whole segment  $\alpha d$ , contrary to Lemma 11.5. That completes the proof that  $e(x) \neq x$ .

Define

$$e_2(a, b, c) := ext(e(a), a, b, c).$$

Then the segment with endpoints  $e(a)$  and  $a$  is not a null segment, so  $e_2(a, b, c)$  is everywhere defined, and if  $d = e_2(a, b, c)$ , we have  $ad = bc$  by (A4). That completes the proof.

**11.3. Hilbert planes and constructive geometry.** In this paper, we have considered line-circle as an axioms. Classically, there is a tradition of studying the consequences of (A1)-(A9) alone, which is known as the theory of Hilbert planes; this theory corresponds to Hilbert's axioms without any form of continuity and without the parallel axiom. The question to be considered here is whether there is an interesting constructive geometry of Hilbert planes. There should be such a theory, with ruler and compass replaced by "Hilbert's tools", which permit one to extend line segments and "transport angles", i.e. to construct a copy of a given angle with specified vertex  $b$  on a specified side of a given line  $L$ . That "tool" corresponds to a Skolem function for Hilbert's axiom C3.

Indeed, most of the development in [19] from A1-A9 is perfectly constructive. In particular, we can prove Hilbert's C3 and the related "triangle construction theorem" enabling us to copy a triangle. But to apply C3, we must have a line  $L$  and a specified side of  $L$ ; to specify a side of  $L$  we must have a point  $p$  not on  $L$ . Consider the proposition that for every line  $L$  there exists a point  $c$  not on  $L$ . In Tarski's language that becomes

$$\forall a, b (a \neq b \rightarrow \exists c (\neg Col(a, b, c))).$$

Classically, the theorem is a trivial consequence of the lower dimension axiom (A8), which gives us three non-collinear points  $\alpha$ ,  $\beta$ , and  $\gamma$ . One of those points will do for  $c$ . But that argument is not constructively valid, since it uses a case distinction to consider whether  $a$  and  $b$  both line on  $Line(\alpha, \beta)$  or not. It is an interesting example, because it illustrates in a simple situation exactly what more is required for a constructive proof than for a non-constructive proof. For a constructive proof, we would need to find a uniform ruler-and-compass construction that applies to any two points  $a$  and  $b$  (determining a line  $L$ ), and produces a point not on  $L$ .

If we could erect a perpendicular to line  $L$  at  $a$ , then (since lines are given by two points) we would already have constructed a point off  $L$ . Gupta constructs perpendiculars without circles: maybe he has solved the problem? No, as it turns out. Gupta's construction has to *start* with a given point  $p$  not on  $L$ . He shows how to construct a perpendicular to  $L$  at  $a$ , but the first step is to draw

the line  $ap$ . The same is true for all the constructions of erected perpendiculars discussed above.

LEMMA 11.7. *Given  $a \neq b$ , there exists a point  $c$  not collinear with  $a$  and  $b$ .*

*Remarks.* We do not know a direct construction; the proof we give uses the introduction of coordinates. It does in principle provide a geometric construction, but it will be very complicated and not visualizable. If we assume circle-circle continuity, we have an easy solution: by the method of Euclid I.1 we produce an equilateral triangle  $abc$ , whose vertex  $c$  can be shown not to lie on line  $L$ . But we note that the proof of circle-circle continuity that we give below via the radical axis requires a point not on the line connecting the centers to get started, so cannot be used to prove this lemma. In order to prove circle-circle continuity without assuming an “extra” point, we also need to introduce coordinates.

*Proof.* Let  $\alpha$ ,  $\beta$ , and  $\gamma$  be the three pairwise non-collinear points guaranteed by (A8). Let  $Line(\alpha, \beta)$  be called the  $x$ -axis. Let 0 be another name for  $\alpha$  and 1 another name for  $\beta$ . Since  $\gamma$  does not lie on the  $x$ -axis, we can use it to erect a perpendicular  $0p$  to the  $x$ -axis at 0, on the other side of the  $x$ -axis from  $\gamma$ . Call that line the  $y$ -axis. Let  $i = ext(p, 0, 0, 1)$ . (Then  $i$  is on the same side of the  $x$ -axis as  $\gamma$ .) Using the uniform perpendicular construction, and the point  $i$  not lying on the  $x$ -axis, we define  $X(p)$  to be the point on the  $x$  axis such that the perpendicular to the  $x$ -axis at  $X(p)$  passes through  $p$ . Similarly we define  $Y(p)$  using the point 1 not on the  $y$ -axis. As shown in [5], using the parallel axiom we can define a point  $p = (x, y)$  given points  $x$  and  $y$  on the  $x$ -axis and  $y$ -axis, such that  $X(p) = x$  and  $Y(p) = y$ , and define addition and multiplication on the  $X$ -axis and prove their field properties. Then coordinate algebra can be used in geometry, and in particular a point not on line  $L$  can be found, by calculating suitable coordinates.

Applying our metatheorems below to this lemma, we see that there is a term  $t(a, b)$  such that  $A(a, b, t(a, b))$  is provable. Of course, since the theorem is classically provable, by Herbrand's theorem there must be a finite number of terms, such that in each case one of those terms will work, and indeed, the three constants  $\alpha$ ,  $\beta$ , and  $\gamma$  illustrate Herbrand's theorem in this case:

$$a \neq b \rightarrow \neg Col(a, b, \alpha) \vee \neg Col(a, b, \beta) \vee \neg Col(a, b, \gamma).$$

Thanks to Lemma 11.7, the classical theory of Hilbert planes has a constructive version, just as the classical theory of ruler-and-compass constructions does.

**§12. Metatheorems.** In this section, we prove some metatheorems about the two Skolemized constructive theories of Tarski geometry, i.e., either intuitionistic Tarski with Skolem functions, or ruler-and-compass Tarski. Both theories have line-circle and circle-continuity with terms for the intersections, and a Skolem function symbol *center* for the triangle circumscription principle; ruler-and-compass Tarski has the logic of partial terms and a symbol  $il(a, b, c, d)$  for the intersection point of two lines, while intuitionistic Tarski with Skolem functions has a Skolem function symbol  $ip$  for inner Pasch. Straightedge and compass constructions correspond to terms of ruler-and-compass Tarski; we have shown that

these can all be imitated by terms of intuitionistic Tarski with Skolem functions, i.e.  $i\ell$  is eliminable.

**12.1. Things proved to exist in constructive geometry can be constructed.** In this section we take up our plan of doing for constructive geometry what cut-elimination and recursive realizability did for intuitionistic arithmetic and analysis, namely, to show that existence proofs lead to programs (or terms) producing the object whose existence is proved. In the case of constructive geometry, we want to produce geometrical constructions, not just recursive constructions (which could already be produced by known techniques, since geometry is interpretable in Heyting’s arithmetic of finite types, using pairs of Cauchy sequences of rational numbers as points).

**THEOREM 12.1** (Geometric constructions extracted from intuitionistic proofs). *Suppose intuitionistic Tarski geometry with Skolem functions proves*

$$P(x) \rightarrow \exists y \phi(x, y)$$

where  $P$  is negative (does not contain  $\exists$  or  $\vee$ ). Then there is a term  $t(x)$  of intuitionistic Tarski geometry with Skolem functions such that

$$P(x) \rightarrow \phi(x, t(x))$$

is provable.

*Proof.* We use cut-elimination. Since our axiomatization is quantifier-free, if  $\psi \rightarrow \exists y \phi$  is provable, then there is a list  $\Gamma$  of quantifier-free axioms such that  $\Gamma, \psi \Rightarrow \exists y \phi$  is provable by a cut-free (hence quantifier-free) proof. Since our axiomatization is disjunction-free, by [14] we can permute the inferences so that the existential quantifier is introduced at the last step. Then we obtain the desired proof just by omitting the last step of the proof. That completes the proof. All the work was in arranging the axiom system to be quantifier-free and disjunction-free.

The term  $t(x)$  in the preceding theorem represents a geometrical construction, but the points constructed by intersecting lines are always given either by *center* or *ip* terms, so the construction contains a “justification” for the fact that the lines intersect. On the other hand, the construction cannot be read literally as a construction script, but requires extra steps to construct the lines implicit in the *center* and *ip* constructions. Moreover, there is nothing in the theorem itself to guarantee that the “definedness conditions” for  $t(x)$  are met, since the Skolem functions are total. The following theorem about ruler-and-compass Tarski geometry does not have that defect, since that theory uses the logic of partial terms.

**THEOREM 12.2** (Geometric constructions extracted from intuitionistic proofs). *Suppose intuitionistic Tarski geometry (without Skolem functions) proves*

$$P(x) \rightarrow \exists y \phi(x, y)$$

where  $P$  is negative (does not contain  $\exists$  or  $\vee$ ). Then there is a term  $t(x)$  of intuitionistic ruler-and-compass Tarski geometry such that

$$P(x) \rightarrow \phi(x, t(x))$$

is also provable. Moreover, if the proof of  $P(x) \rightarrow \exists y \phi(x, y)$  does not use certain axioms, then the term  $t(x)$  does not involve the Skolem symbols for the unused axioms.

*Proof.* We have a choice of two proofs. We could use cut-elimination directly, but then we need it for the logic of partial terms and not just for ordinary intuitionistic predicate calculus. The details of the cut-elimination theorem for such logics have not been published, but they are not significantly different from Gentzen's formulation for first-order logic. While we are explaining this point, it is no more complicated to explain it for multi-sorted theories with LPT, which were used in [3] with axioms for Hilbert-style geometry. Specifically, we reduce such theories to ordinary predicate calculus as follows: introduce a unary predicate for each sort, and then if  $t$  is a term of sort  $P$ , interpret  $t \downarrow$  as  $P(t)$ . Now we have a theory in first-order one-sorted predicate calculus, which is quantifier-free and disjunction-free if the original theory was, and we can apply ordinary cut-elimination, as in the proof of Theorem 12.1.

Alternately, we can avoid using cut-elimination for LPT, by first translating the original formula from ruler-and-compass Tarski to intuitionistic Tarski with Skolem functions using Theorem 10.5. Then the resulting construction term  $t$  involves the function symbol  $ip$ ; but it is easy to express  $ip$  in terms of  $il$  if that is desired, i.e. to interpret intuitionistic Tarski with Skolem functions into ruler-and-compass Tarski. That completes the proof.

**12.2. Extracting constructions from classical proofs.** The following theorem illustrates the essential difference between constructive and classical (non-constructive) geometry: in a constructive existence theorem, we must supply a single (uniform) construction of the point(s) whose existence is asserted, but in a classical theorem, there can be several cases, with a different construction in each case.

**THEOREM 12.3** (Constructions extracted from classical proofs). *Suppose classical Tarski geometry with Skolem functions proves*

$$P(x) \rightarrow \exists y \phi(x, y)$$

where  $P$  is quantifier-free and disjunction-free. Then there are terms  $t_i(x)$  such that

$$P(x) \rightarrow \phi(x, t_1(x)) \vee \dots \vee \phi(x, t_n(x))$$

is also provable.

*Proof.* This is a special case of Herbrand's theorem.

*Example 1.* There exists a perpendicular to line  $L$  through point  $p$ . Classically, one argues by cases: if  $p$  is on  $L$ , then we can "erect" the perpendicular, and if  $p$  is not on  $L$  then we can "drop" the perpendicular. So the proof provides two constructions,  $t_1$  and  $t_2$ . This is not a constructive proof. In [5], we give a more complicated uniform construction. When adapted to Tarski's system, this constructive gives a single term to construct a perpendicular.

*Example 2.* Euclid's proof of Book I, Proposition 2 provides us with two such constructions,  $t_1(a, b, c) = c$  and  $t_2(a, b, c)$  the result of Euclid's construction of

a point  $d$  with  $ad = bc$ , valid if  $a \neq b$ . Classically we have  $\forall a, b, c \exists d(ad = bc)$ , but we need two terms  $t_1$  and  $t_2$  to cover all cases.

*Example 3.* Let  $p$  and  $q$  be distinct points and  $L$  a given line, and  $a, b$ , and  $c$  points on  $L$ , with  $a$  and  $b$  on the same side of  $L$  as  $c$ . Then there exists a point  $d$  which is equal to  $p$  if  $b$  is between  $a$  and  $c$  and equal to  $q$  if  $a$  is between  $b$  and  $c$ . The two terms  $t_1$  and  $t_2$  for this example can be taken to be the variables  $p$  and  $q$ . One term will not suffice, since  $d$  cannot depend continuously on  $a$  and  $b$ , but all constructed points do depend continuously on their parameters. This classical theorem is therefore not constructively provable.

**12.3. Disjunction properties.** We mentioned above that intuitionistic Tarski geometry cannot prove any non-trivial disjunctive theorem. That is a simple consequence of the fact that its axioms contain no disjunction. We now spell this out:

**THEOREM 12.4** (No nontrivial disjunctive theorems). *Suppose intuitionistic Tarski geometry proves  $H(x) \rightarrow P(x) \vee Q(x)$ , where  $H$  is negative. Then either  $H(x) \rightarrow P(x)$  or  $H(x) \rightarrow Q(x)$  is also provable. (This result depends only on the lack of disjunction in the axioms.)*

*Proof.* Consider a cut-free proof of  $\Gamma, H(x) \rightarrow P(x) \vee Q(x)$ , where  $\Gamma$  is a list of some axioms. Tracing the disjunction upwards in the proof, if we reach a place where the disjunction was introduced on the right before reaching a leaf of the proof tree, then we can erase the other disjunct below that introduction, obtaining a proof of one disjunct as required. If we reach a leaf of the proof tree with  $P(x) \vee Q(x)$  still present on the right, then it occurs on the left, where it appears positively. Its descendants will also be positive, so it cannot participate in application of the rule for proof by cases (which introduces  $\vee$  in the left side of a sequent); and it cannot reach left side of the bottom sequent, namely  $\Gamma, H(x)$ , as these formulas contain no disjunction. But a glance at the rules of cut-free proof, e.g. on p. 442 of [14], will show that these are the only possibilities. That completes the proof.

We note that order on a fixed line  $L$  can be defined using betweenness, so it makes sense to discuss the provability of statements about order.

**COROLLARY 12.5.** *Intuitionistic Tarski geometry does not prove apartness*

$$a < b \rightarrow x < b \vee a < x.$$

*Proof.* The statement in question is a disjunctive theorem, so the theorem applies.

**COROLLARY 12.6.** *Intuitionistic Tarski geometry does not prove the principle  $x \neq 0 \rightarrow x < 0 \vee x > 0$  or the equivalent principle that if point  $p$  does not lie on line  $L$ , then any other point  $x$  is either on the same side of  $L$  as  $p$  or the other side.*

*Proof.* The statement in question is a disjunction theorem, so the theorem applies.

**12.4. Interpretation of Euclidean field theory.** A Euclidean field is defined constructively as an ordered ring in which nonzero elements have reciprocals. The relation  $a < b$  is primitive;  $a \leq b$  abbreviates  $\neg b < a$ . The axioms of Euclidean field theory include stability of equality and order. Stability of order, that is  $\neg b \leq a \rightarrow a < b$ , is also known as Markov's principle. Classically, the models of ruler and compass geometry are planes over Euclidean fields. We showed in [5] that a plane over a Euclidean field is a model of ruler-and-compass geometry, when ruler-and-compass geometry is defined in any sensible way; constructively, this theorem takes the form of an interpretation  $\phi \mapsto \bar{\phi}$  from some geometric formal theory to the theory **EF** of Euclidean fields.

The converse direction is much more difficult; we have to show that any model of geometry is a plane over a Euclidean field  $F$ . To do that, we fix a line  $F$  to serve as the  $x$ -axis (and the domain of the field); fix a point  $0$  on that line, erect a perpendicular  $Y$  to  $F$  at  $0$  to serve as the  $y$ -axis. Given any pair of points  $(x, y)$  on  $F$ , we rotate  $y$  by ninety degrees to a point  $y'$  on the  $y$ -axis, and then erect perpendiculars at  $x$  to  $F$  and at  $y'$  to  $Y$ . These perpendiculars should meet at a point  $MakePoint(x, y)$ . It is possible to show by the strong parallel axiom that they do meet. This construction is the starting point for the following theorem:

**THEOREM 12.7.** *Every model of intuitionistic Tarski geometry is a plane over a Euclidean field. Moreover, there is an interpretation  $\phi \mapsto \phi^\circ$  from the theory of Euclidean fields to intuitionistic Tarski geometry.*

*Proof.* In addition to introducing coordinates as discussed above, one also has to define addition and multiplication geometrically in order to interpret the addition and multiplication symbols of Euclidean field theory. It has been shown in [5] how to do this; the proofs there can be formalized in intuitionistic Tarski geometry, so we obtain a model-theoretic characterization of the models of that theory.

Moreover, our work with the double-negation interpretation above can now be put to good use. For example, the definition of multiplication can be given directly following Hilbert's definition, which is based on the triangle circumscription principle. It is easy to give a term  $HilbertMultiply(a, b)$  that takes two points  $a$  and  $b$  on a fixed line (the " $x$ -axis") and produces their product (also a point on the  $x$ -axis), using *center* and the uniform rotation construction. (See [5] for details.) But once that term is given, the assertions that it satisfies the associative and commutative laws are quantifier-free, and hence, the proofs in [19] are "importable." Technically, one must check that the degenerate cases of inner Pasch are not used, but that is all that one has to check by hand. In [5], there is a definition of "uniform addition", i.e. without a case distinction on the signs of the addends. A term  $Add(x, y)$  defining the sum of  $x$  and  $y$  is given in [5]. Again, once the term is given, we can be assured by the double-negation interpretation that its properties are provable in intuitionistic Tarski with Skolem functions, if we just check [19] to make sure the degenerate cases of inner Pasch are not used.

The terms  $Add$  and  $HilbertMultiply$  can then be used to define a syntactic interpretation  $\phi \mapsto \phi^\circ$  from the theory of Euclidean fields to intuitionistic Tarski geometry. That completes the proof of the theorem.

**§13. Circle-circle continuity.** In this section we show that circle-circle continuity is a theorem of intuitionistic Tarski geometry; that is, we can derive the existence of the intersection points of two circles (under the appropriate hypotheses). The similar theorem for classical Tarski geometry can be derived indirectly, using the representation theorem (Theorem 9.7) and Gödel’s completeness theorem; but for intuitionistic Tarski geometry, we must actually exhibit a construction for the intersection points of two circles, and prove constructively that it works. This question relies on Euclid III.35, a theorem about how two chords of a circle divide each other into proportional segments, and III.36, a similar theorem, and constructively it requires a combined version of those two propositions without a case distinction as to which applies (i.e., two lines cross inside or outside a circle). Aside from those theorems, it uses only very straightforward geometry.

**THEOREM 13.1.** *In intuitionistic Tarski geometry with only line-circle continuity: given two circles  $C$  and  $K$  satisfying the hypotheses of circle-circle continuity, and a point not on the line  $L$  connecting their centers, we can construct the point(s) of intersection of  $C$  and  $K$ .*

There are two approaches to proving this theorem, which we will discuss separately. The first method proceeds by introducing coordinates and reducing the problem to algebraic calculations by analytic geometry. While in principle this does produce an ultimately purely geometric proof, one cannot visualize the lengthy sequence of constructions required. For esthetic reasons, therefore, we also give a second proof, which avoids the use of coordinates by the use of a well-known construction called the “radical axis.” In this more geometric proof, essential use is made of the “extra point” in the hypothesis. That point can of course be constructed, by Lemma 11.7, but to do so we have to introduce coordinates, which we wish to avoid here for esthetic reasons.<sup>22</sup>

**13.1. Circle-circle continuity via analytic geometry.** Given two circles  $C$  and  $K$  with distinct centers  $s$  and  $t$ , let  $L$  be the line through the centers. Given a point not on  $L$ , we can erect a perpendicular to  $L$  at  $s$ , and introduce coordinates in the manner of Descartes and Hilbert, with constructive extension to negative arguments as developed in [5]. We can choose the point  $t$  as 1. Now the tools of analytic geometry are available. Let  $r$  be the radius of circle  $C$  and  $R$  the radius of circle  $K$ , and calculate the equations of  $C$  and  $K$  and solve for a point  $(x, y)$  lying on both circles. It turns out that some crucial terms cancel, and we can solve the equations using only square roots, which means that we can solve them geometrically using the methods of Descartes and Hilbert, with the constructive modifications *op. cit.* To derive circle-circle continuity, we must show that the hypothesis that circle  $C$  has a point inside circle  $K$  makes the quantities under the square root non-negative, and the extra hypothesis that the circle  $C$  has a point strictly inside  $K$  makes the two solutions of the equation distinct.

---

<sup>22</sup>Both these proofs use the parallel axiom essentially. There is a proof in the literature that line-circle continuity implies circle-circle continuity without the use of the parallel axiom [22]. It is based on Hilbert’s axioms rather than Tarski’s, and we do not know if it is constructively justifiable or not. But we have no reason to avoid the parallel axiom for present purposes.

A similar theorem is proved classically in [12], p. 144, but a few details are missing there. The issue is that it is not enough to observe that the equations for the intersection points are quadratic. One has to translate the hypothesis that one circle has a point inside and a point outside the other circle into algebra and show algebraically that this implies the equations for the intersection are solvable.

This is a fairly routine exercise and is all perfectly constructive; but it is somewhat unsatisfying to have to resort to coordinates. One would like to see a direct geometric construction of the points of intersection of two circles, using only a few steps, rather than the dozens or perhaps hundreds of not-visualizable steps required to geometrize an algebraic calculation. There is indeed such a geometric construction, using the “radical axis” of the two circles. Below we verify that the radical axis construction can be carried out constructively (i.e. does not require any case distinctions in its definition), and that the correctness proof can be carried out in intuitionistic Tarski geometry. Although the construction itself is easy to visualize (it is only a few steps with ruler and compass), the correctness proof in Tarski geometry is more complicated.

**13.2. Euclid Book III in Tarski geometry.** The correctness proof of the radical axis construction requires the last two propositions of Euclid Book III; and moreover its formalization in Tarski geometry is of independent interest.

Book III of Euclid can be formalized in intuitionistic Tarski geometry, but since most of the theorems mention angles, we need to use the developments of Chapter 11 of [19], where angles, angle congruence, etc. are developed. However, some propositions can be proved quite simply, for example III.31 (an angle inscribed in a semicircle is right), which goes back to I.29 and I.11 and hence to the construction of perpendiculars (and not to Book II at all). We also will need a related proposition that might have been (but does not seem to be) in Euclid:

LEMMA 13.2. (*in Tarski geometry with segment-circle continuity*) *If  $axb$  is a right angle and  $ab$  is a diameter of a circle  $C$  then  $x$  lies on  $C$ .*

*Proof.* By segment-circle continuity, we can find a point  $y$  on  $C$  and on the ray from  $a$  through  $x$ . Then by Euclid III.30,  $ayb$  is a right angle, so if  $y \neq x$  then  $xb$  and  $yb$  are two perpendiculars to  $ay$  through  $b$ , contradiction. By the stability of equality we have  $y = x$ , so  $x$  lies on  $C$ . That completes the proof.

We present here another proposition from Euclid Book III that can be proved directly from the Tarski axioms.

LEMMA 13.3 (Euclid III.18). *Suppose line  $L$  meets circle  $C$  with center  $c$  in exactly one point  $a$ . Then  $ca \perp L$ .*

*Remark.* We need dropped perpendiculars to prove this lemma, but we already derived the existence of dropped perpendiculars from line-circle continuity, so that is not a problem. The proof uses Euclid’s idea, but Tarski’s definition of perpendicular.

*Proof.* Drop a perpendicular  $cb$  from  $c$  to line  $L$ , which can be done by Lemma 7.5. We want to prove  $b = a$ . By the stability of equality, we can proceed by contradiction, so suppose  $b \neq a$ . Point  $b$  is outside the circle (i.e. there is a point  $e$  on  $C$  with  $\mathbf{B}(c, e, b)$ , since otherwise by line-circle continuity,  $L$  meets  $C$  in a

second point. Let  $e$  be the reflection of  $a$  in  $b$ ; then  $ab = be$ , and since  $ce \perp L$ , we have  $ca = ce$ . That is,  $e$  lies on circle  $C$  as well as line  $L$ , contradicting the hypothesis that  $L$  meets  $C$  only once. That completes the proof.

Despite these examples, the radical axis construction that we use makes use of some developments of Euclid Book III that are not so straightforward, because they rest on the theory of proportionality in Euclid Book II. We want to make sense of the phrase

$$ab \cdot cd = pq \cdot rs.$$

In Euclid, this is written “the rectangle contained by  $ab$  and  $cd$  is equal to the rectangle contained by  $pq$  and  $rs$ .” But Euclid has no concept of “area” as represented by (the length of) a segment. To define this relation geometrically, we could use a definition of similar triangles, involving two right triangles with sides  $ab$ ,  $pq$  and  $cd$ ,  $rs$  respectively. Book II of Euclid develops (quadratic) algebra on that basis.

There is another way to define the notion  $ab \cdot cd = pq \cdot rs$ . Namely, introduce coordinates, define multiplication geometrically, and interpret  $ab \cdot cd$  as multiplication of segments on the  $x$ -axis congruent to  $ab$  and  $cd$ . This is not actually so different from the first (Euclidean) interpretation, since similar triangles are used in defining multiplication. For the modern analysis of Euclid’s notion of equality for “figures” see page 197 of [12], and for the connection to geometric multiplication see page 206. There it is proved that equality in Euclid’s sense corresponds to algebraic equality using geometric arithmetic; in particular, the two definitions of  $ab \cdot cd = pq \cdot rs$  are provably equivalent.

To carry out the radical axis construction constructively, we need to extend the notion  $ab \cdot cd$  to allow signed segments. To do this directly using similar triangles would be to duplicate the effort of defining signed multiplication in [5]. Therefore we use the geometric-multiplication definition, following [12].


Either definition can be expressed in a quantifier-free way using intuitionistic Tarski geometry with Skolem functions, so the two notions are provably equivalent in intuitionistic Tarski geometry if and only if they are provably equivalent in classical Tarski geometry. Since Hartshorne *op. cit.* proves them equivalent in classical Hilbert geometry (and without using circle-circle continuity), and since Hilbert’s axioms (except continuity) are provable in classical Tarski geometry (without continuity) (as shown in [19]), it follows that the two definitions are provably equivalent in intuitionistic Tarski geometry.

We need to define the notion of the power of a point in intuitionistic Tarski geometry, and check that its principal properties can be proved there. That notion is usually defined as follows, when  $b$  is not the center of  $C$ : Let  $c$  be the center of  $C$ , and  $x$  the point of intersection of  $\text{Line}(b, c)$  with  $C$  that is on the same side of  $c$  as  $b$ , and  $y$  the other point of intersection. Then the power of  $b$  with respect to  $C$  is  $bx \cdot by$ . If we interpret the dot as (signed) multiplication (see [5] of (directed) segments on the  $x$ -axis of points congruent to the segments mentioned, then this definition makes sense in intuitionistic Tarski geometry. It does, however, have the disadvantage that the power of the center of the circle is not defined; and we cannot just define it to be  $-1$ , as we could classically, because constructively we cannot make the case distinction whether  $b$  is or is

not the center. This definition can be fixed constructively as follows. Fix a diameter  $pq$  of circle  $C$ , whose center is  $c$ . Then given any point  $b$ , extend segment  $pc$  by  $bc$  to produce point  $B$ . (If  $b = c$ , this is still legal and produces  $c$ .) Then the power of  $b$  with respect to  $C$  is  $Bq \cdot Bp$ , where the dot is signed multiplication (so that  $Bq$  and  $Bp$  have opposite signs when  $b$  is inside  $C$ .) This gives the same answer as the usual definition when  $b$  is not the center.

The following lemma shows that the power of  $p$  with respect to  $C$  can be computed from any chord, not just from the diameter. See Fig. 25.

FIGURE 25. The power of  $b$  with respect to  $c$  can be computed from any chord, because  $ba \cdot bc = bd \cdot be$ .



LEMMA 13.4. *In Tarski geometry with only line-circle continuity: Let  $C$  be a circle with center  $c$  and let  $b$  be any point. Let  $L$  be any line through  $b$  meeting  $C$  in points  $u$  and  $v$ . Then the power of  $C$  with respect to  $C$  is  $ub \cdot vb$ .*


*Remark.* For the case when  $b$  is inside  $C$  (first part of Fig. 25), this is Euclid III.35, and for  $b$  outside it is III.36. The general case is mentioned in Heath's commentary on III.35, as a corollary of III.35 and III.36. It also occurs as Exercise 20.3 in [12]. The proof implicitly suggested there by Exercise 20.2 is the same one suggested in Heath's commentary.

*Proof.* Hilbert multiplication can be defined by Skolem terms in intuitionistic Tarski geometry, without circle-circle continuity, as the constructions in [5] show. To recap: multiplication is defined by using the triangle circumscription axiom to draw a circle, and then the product is given by the intersection of that circle with a line. In addition there are some rotations involved, which also can be defined by terms. When formulated in intuitionistic Tarski geometry with Skolem functions, the statement of the lemma is quantifier-free. By the double-negation interpretation, it is provable constructively if and only if it is classically provable. And it is classically provable, cf. the exercise mentioned in the remark. Of course, the textbook containing the exercise is based on Hilbert's axioms, but [19] derives all of Hilbert's axioms from Tarski's (classically), so once you solve Exercise 20.3 *op. cit.*, it follows that the result is provable in intuitionistic Tarski geometry with Skolem functions.

**13.3. The radical axis.** In this section, we discuss the construction of the "radical axis" of two circles, with attention to constructivity. In the next section we will use the radical axis to give a second proof that line-circle continuity implies circle-circle continuity.

The “radical axis” of two circles is a line, defined whether or not the circles intersect, such that if they do intersect, the line passes through the points of intersection (and if they are tangent, it is the common tangent line). On page 182 of [12], a ruler-and-compass construction of the radical axis is given. Fig. 26 illustrates the construction, for the benefit of readers who do not have [12] at hand.<sup>23</sup>

FIGURE 26. Construction of the radical axis.  
 $m = \text{midpoint}(a, b)$ ; draw  $sm$  and  $mt$  and drop perpendiculars from  $a$  to  $sm$  and from  $b$  to  $mt$ . Their intersection is  $p$  and the radical axis is perpendicular to  $st$  through  $p$ .



The initial data are the centers  $s$  and  $t$  of the two circles, with  $s \neq t$ , and two points  $a$  and  $b$  on the circles, such that  $a \neq b$  and  $ab$  does not meet the line joining the centers. (That hypothesis allows one of the circles to be a null circle (zero radius), but not both). The construction is as follows: First define  $m$  as the midpoint of  $ab$ . Then  $\mathbf{B}(a, m, b)$ . If  $s = m$  or  $t = m$  then  $ab$  meets  $st$ , contrary to hypothesis. Hence we can construct the lines  $sm$  and  $tm$ . Then construct perpendiculars to those lines through  $a$  and  $b$  respectively (using the uniform perpendicular, so we do not need to worry if  $a$  and  $b$  are on those lines or not).

<sup>23</sup>The radical axis was already old in 1826 [20], although there it is constructed from the intersection points of circles, rather than the other way around. I do not know the origin of the ruler-and-compass construction used here.

Then  $p$  is to be the intersection of these two perpendiculars, and the radical axis is the perpendicular to  $st$  through  $p$ , again using the uniform perpendicular.

**LEMMA 13.5.** *Given two circles  $C$  and  $K$ , the radical axis as constructed above does not depend on the particular points  $a$  and  $b$  chosen, and can be constructed from the two circles and one additional point not lying on the line joining the centers.*

*Remark.* The “extra point” is a necessary parameter. The circles are presumed given by center and point, but the points giving the circles might happen to lie on the center line.

*Proof.* Let the two circles  $C$  and  $K$  have centers  $s$  and  $t$ , and let  $a$  and  $b$  be distinct points on  $C$  and  $K$ , respectively, such that the midpoint  $m$  of  $ab$  does not lie on the line  $L$  containing the centers  $s$  and  $t$ . We need a point  $r$  not on  $L$  to be able to choose  $a$  and  $b$ . For example, we can use  $r$  to erect a perpendicular to  $L$  at  $s$ , and let  $a$  be one of its intersections with  $C$ , and then construct a perpendicular to  $L$  at  $t$ , on the same side of  $L$  as  $a$ , and let  $b$  be the intersection of this perpendicular with  $K$ .

We wish to construct the radical axis  $R$  of  $C$  and  $K$ . In implementing the construction given above in intuitionistic Tarski geometry, we must appeal to Lemma 7.9 for the construction of midpoints using only line-circle continuity, and to the uniform perpendicular construction to construct the required perpendiculars, so that we do not need to worry about whether  $a$  lies on  $sb$  or  $b$  lies on  $sa$ . But we do have to worry about whether the intersection  $p$  of those perpendiculars exists. By the strong parallel axiom, it will exist if the two perpendiculars are not parallel or coincident. That can only happen if  $m$ ,  $s$ , and  $t$  are collinear, but we have chosen  $a$  and  $b$  so that  $b$  does not lie on  $L$ ; hence indeed  $p$  exists. Then define line  $R$  as the (uniform) perpendicular to  $L$  through  $p$ .


Now we will prove (constructively and using only line-circle continuity) that every point  $x$  on  $R$  has equal powers with respect to  $C$  and  $K$ . Suppose  $x$  is on the radical axis  $R$ . Define circle  $M$  to be the circle with center  $m$  and passing through  $a$  and  $b$ . See Fig. 27.

Let  $z$  be the intersection of  $M$  with  $\text{Line}(a, p)$ , and let  $y$  be the intersection of  $M$  with  $\text{Line}(b, x)$ . Then the power of  $x$  with respect to  $C$ , namely  $xz \cdot ax$  (by Lemma 13.4) is equal to the power of  $x$  with respect to  $M$ . Similarly, the power of  $x$  with respect to  $K$  is  $yx \cdot yb$ , which is also the power of  $x$  with respect to  $M$ . Since the powers of  $x$  with respect to  $C$  and to  $K$  are both equal to the power of  $x$  with respect to  $M$ , they are equal. That completes the proof.

**13.4. Circle-circle continuity via the radical axis.** We observe that the power of  $x$  with respect to circle  $C$  is negative when  $x$  is inside  $C$ , zero when  $x$  is on  $C$ , and positive when  $x$  is outside  $C$ . Intuitively, the radical axis of two intersecting circles is the line joining the two intersection points. In this section we use this idea to prove circle-circle continuity. One more idea is necessary: the “radical center.”

*Second proof of Theorem 13.1.* Let circles  $C$  and  $K$  be given, satisfying the hypothesis of the circle-circle continuity axiom. That is,  $K$  has a point (say  $b$ )

FIGURE 27. The power of  $p$  with respect to each of  $C$  and  $K$  is equal to the power of  $p$  with respect to  $M$ .



outside  $C$  and a point (say  $x$ ) inside  $C$ . We will use  $b$  as one of the points to start the radical axis construction, and we will choose the other point  $a$  very carefully. Construct the perpendicular to  $xb$  at  $x$ . Since  $x$  is inside  $C$ , by segment-circle continuity that perpendicular meets  $C$  in some point  $a$  such that the interior of  $ab$  does not meet  $Line(s, t)$ . In other words,  $a$  and  $b$  lie on the same side of  $Line(s, t)$  unless  $b$  already lies on  $Line(s, t)$ .

Now use  $a$  and  $b$  as starting points for the radical axis construction; let the constructed point be  $p$ . By construction of  $a$ , if  $x \neq b$ , then angle  $axb$  is a right angle with  $a$  and  $x$  at the ends of a diameter of circle  $M$ ; that is the result of choosing  $a$  as we did. See Fig. 28. Therefore the vertex  $x$  of this right angle lies on circle  $M$ , by Lemma 13.2. On the other hand, if  $x = b$  (that is  $xb$  is tangent to  $K$ ) then  $x$  also lies on  $M$ ; by the stability of equality,  $x$  lies on  $M$  (that is  $xt = bt$ ) whether or not  $x = b$ .

By segment-circle continuity, there is a point  $v$  on  $C$  with  $\mathbf{B}(v, x, p)$ . We claim  $\mathbf{T}(v, p, b)$ . By the stability of betweenness we can prove this by contradiction. Since  $\neg\neg(A \vee B)$  is equivalent to  $\neg(\neg A \wedge \neg B)$ , we can argue by cases for the contradiction. There are two cases to consider:  $\mathbf{B}(p, v, b)$  and  $\mathbf{B}(v, b, p)$ .



$x$  lies on  $R$ , the power of  $x$  with respect to  $K$  is equal to the power of  $x$  with respect to  $C$ , which is zero. Hence  $x$  lies on  $K$  as well as on  $C$ . That completes the proof.

We have shown that points on the radical axis have equal powers with respect to both circles. The following lemma is the converse. We do not need it, but the proof is short and pretty.

**LEMMA 13.6.** *Let  $C$  and  $K$  be distinct circles. If they meet, then the radical axis of  $C$  and  $K$  consists of exactly those points whose powers with respect to  $C$  and  $K$  are equal.*

*Remark.* The lemma is true even if the circles do not meet, but I do not know a simple geometric proof.

*Proof.* We have already proved that points on the radical axis have equal powers. It suffices to prove the converse. Suppose that  $u$  has equal powers with respect to  $C$  and  $K$ . We must prove  $u$  lies on the radical axis. Let  $v$  lie on both circles. When we compute the powers of  $u$  with respect to both  $C$  and  $K$  using the line  $uv$ , we get different answers unless  $u$  lies on the radical axis (so that the endpoints on  $C$  and  $K$  are the same). That completes the proof.

**13.5. Skolem functions for circle-circle continuity.** Terms of Tarski geometry (intuitionistic or continuous, which has classical logic) correspond to (certain) ruler-and-compass constructions; in effect, to constructions in which you can form the intersection of lines that must intersect by inner Pasch, and intersections of lines and circles. Since inner Pasch implies outer Pasch, the points formed by outer Pasch are also given by terms; and since circle-circle continuity is implied by line-circle continuity, then there must be terms for constructing those intersection points as well.

But there is an issue to consider, in that the terms for erected and uniform perpendiculars have an extra parameter, a point not on the line; and the radical axis construction also has an extra parameter for a point not on the center line. Examination of those constructions reveals that if the “extra” point is changed to the other side of the line, then the “head” of the perpendicular changes sides too; and in the radical axis construction, the result is that the two intersection points of the two circles switch places. If we then fix that choice once and for all by using Lemma 10.4, we can construct terms that give the two intersection points of two circles continuously. However, those terms will be complicated, because the term for constructing a point not on a line involves coordinates and cross products.

In previous work on constructive geometry, we had circle-circle continuity as an axiom, and built-in function symbols for the intersection points. It was a point of difficulty to distinguish the two, which we wanted to do by saying whether the triple of points from the center of one circle to the other center to the point of intersection was a “right turn” or a “left turn”. The concepts  $Right(a, b, c)$  and  $Left(a, b, c)$  had to be defined, either by a complicated set of axioms, or by introducing coordinates and using cross products as in [5]. Here we can recover those same terms, but now the coordinates and cross products are at least no

longer in the axioms! If we perform a complex conjugation (i.e. reflect in the  $x$ -axis) then the two terms for the intersection points of two circles change places, exactly as in [5].

If one wishes (for example for connecting these theories to computer graphics) to have explicit function symbols for circle-circle continuity, of course they can be conservatively added.

**§14. Conclusion.** We have exhibited a constructive version of Tarski's Euclidean geometry. Because of the double-negation interpretation, it can prove at least some version of each classical theorem. Using the uniform perpendicular, rotation, and reflection constructions given in this paper, it is possible (by the methods of [5]) to give geometric definitions of addition and multiplication, without case distinctions as to the sign of the arguments, and proofs of their properties, so that coordinates in a Euclidean field provably exist. Hence the theory has not omitted anything essential. To achieve these results, we had to modify Tarski's axioms to eliminate degenerate cases, and add back some former axioms that Tarski had eliminated using those degenerate cases. Even with classical logic, this theory now connects nicely with ruler-and-compass constructions, since each of the points asserted to exist can be constructed with ruler and compass.

By cut-elimination, things proved to exist (under a negative hypothesis, as is always the case in Euclid) can be constructed, by a uniform straightedge-and-compass construction. Even stronger, these constructions need not involve taking the intersections of arbitrary lines, but only those lines that have to intersect by the strong parallel axiom or inner Pasch.

By contrast, in Tarski's (classical) theory, we obtain (by Herbrand's theorem) a similar result but without uniformity, i.e., there are several constructions (not necessarily just one), such that for every choice of the "given points", one of the constructions will work. (The classical result (unlike the constructive one) holds only for formulas  $\forall x \exists y A(x, y)$ , where  $A$  is quantifier-free.)

These points-only axiom system have conservative extensions with variables for lines and circles, and further conservative extensions with variables for angles, segments, and arcs, which can serve for the direct constructive formalization of Euclidean geometry using Hilbert's primitives (as in [3]). Therefore, this points-only theory, with its short list of axioms, can be said to provide the logical foundations of constructive Euclidean geometry. In particular, it supplies one detailed example of a formalization of constructive geometry, to which the independence results about the parallel postulate of [5] apply.

**§15. Listing of axioms for reference.** In the following,  $ab = cd$  abbreviates  $E(a, b, c, d)$ , and  $\mathbf{T}(a, b, c)$  is non-strict betweenness, while  $\mathbf{B}(a, b, c)$  is strict betweenness.

**15.1. Classical two-dimensional Tarski geometry.** We give the version preferred by Szmielew. The version in [19] has (the classically equivalent) (A10) instead of (A10<sub>3</sub>). We also give the Skolemized versions here.  $Col(a, b, c)$  (collinearity) is an abbreviation for  $\mathbf{T}(a, b, c) \vee \mathbf{T}(b, c, a) \vee \mathbf{T}(c, a, b)$ .

$ab = ba$	(A1) Reflexivity of equidistance
$ab = pq \wedge ab = rs \rightarrow pq = rs$	(A2) Transitivity of equidistance
$ab = cc \rightarrow a = b$	(A3) Identity of equidistance
$\exists x (\mathbf{T}(q, a, x) \wedge ax = bc)$	(A4) Segment extension
$\mathbf{T}(q, a, ext(q, a, b, c)) \wedge E(q, ext(q, a, b, c), b, c)$	(A4), Skolemized
$(a \neq b \wedge \mathbf{T}(a, b, c) \wedge \mathbf{T}(A, B, C) \wedge ab = AB \wedge bc = BC$ $ad = AD \wedge bd = BD) \rightarrow cd = CD$	(A5) Five-segment axiom
$\mathbf{T}(a, b, a) \rightarrow a = b$	(A6) Identity for betweenness
$\mathbf{T}(a, p, c) \wedge \mathbf{T}(b, q, c) \rightarrow \exists x (\mathbf{T}(p, x, b) \wedge \mathbf{T}(q, x, a))$	(A7) inner Pasch
$\mathbf{T}(a, p, c) \wedge \mathbf{T}(b, q, c) \rightarrow$ $\mathbf{T}(p, ip(a, p, c, b, q), b) \wedge \mathbf{T}(q, ip(a, p, c, b, q), a)$	(A7), Skolemized
$\exists a, b, c \neg Col(a, b, c)$	(A8), lower dimension
$\neg Col(\alpha, \beta, \gamma)$	(A8), Skolemized
$pa = pb \wedge qa = qb \wedge ra = rb \rightarrow Col(a, b, c)$	(A9), upper dimension
$\neg Col(a, b, c) \rightarrow \exists x (ax = bx \wedge ax = cx)$	(A10 <sub>3</sub> ), triangle circumscription
$ax = ap \wedge \mathbf{T}(a, x, b) \wedge \mathbf{T}(a, b, y) \wedge ay = aq \rightarrow$ $\exists z (\mathbf{T}(p, z, q) \wedge az = ab)$	segment-circle continuity

In the Skolemized version of the triangle circumscription principle,  $x$  is given by  $center(a, b, c)$ . We make no use of a Skolemized version of segment-circle, so we do not give one.

**15.2. Intuitionistic Tarski geometry.** This theory takes  $\mathbf{B}$  as primitive rather than  $\mathbf{T}$ , so  $\mathbf{T}(a, b, c)$  is an abbreviation for  $\neg(a \neq b \wedge b \neq c \wedge \neg\mathbf{B}(a, b, c))$ , and  $Col(a, b, c)$  is an abbreviation for

$$a \neq b \wedge \neg(\neg\mathbf{B}(p, a, b) \wedge \neg\mathbf{B}(a, p, b) \wedge \neg\mathbf{B}(a, b, p) \wedge a \neq p \wedge b \neq p),$$

which is equivalent to the double negation of the classical definition of  $Col(a, b, c)$  together with  $a \neq b$ . In other words,  $Col(a, b, c)$  says  $c$  lies on  $Line(a, b)$ . The axioms (A1)-(A3) and (A5) are unchanged, except that now  $\mathbf{T}$  is defined in terms of  $\mathbf{B}$ . It is inessential whether  $\mathbf{T}$  or  $\mathbf{B}$  is taken as primitive.

The differences between classical and intuitionistic Tarski geometry are

- (A4): Only non-null segments can be extended.
- inner Pasch (A7): The hypothesis  $a \neq p \wedge b \neq q \wedge \neg Col(a, b, c)$  is added.
- Symmetry and inner transitivity of betweenness (A14) and (A15) are added.
- A negative formula is used for collinearity in the dimension axioms and the triangle circumscription principle.
- In line-circle continuity, the two points  $p$  and  $q$  determining the line are assumed to be unequal, and we use two-point line-circle continuity instead of segment-circle.
- We use intuitionistic logic and add the stability axioms.

Intuitionistic Tarski geometry plus classical logic is called “continuous Tarski geometry”; we can have continuous Tarski geometry with or without Skolem functions. The changed axioms are as follows:

$$\begin{aligned}
q \neq a &\rightarrow \exists x (\mathbf{T}(q, a, x) \wedge ax = bc) && \text{(A4-i) Segment extension} \\
q \neq a &\rightarrow \mathbf{T}(q, a, \text{ext}(q, a, b, c)) \wedge E(q, \text{ext}(q, a, b, c), b, c) && \text{(A4-i), Skolemized} \\
&\neg \mathbf{B}(a, b, a) && \text{(A6-i)} \\
\mathbf{T}(a, p, c) \wedge \mathbf{T}(b, q, c) \wedge p \neq a \wedge p \neq c \wedge \neg \text{Col}(a, b, c) &\rightarrow \\
\exists x (\mathbf{B}(p, x, b) \wedge \mathbf{B}(q, x, a)) &&& \text{(A7-i) strict inner Pasch} \\
\mathbf{T}(a, p, c) \wedge \mathbf{T}(b, q, c) \wedge p \neq a \wedge p \neq c \wedge \neg \text{Col}(a, b, c) &\rightarrow \\
\mathbf{B}(p, \text{ip}(a, p, c, b, q), b) \wedge \mathbf{B}(q, \text{ip}(a, p, c, b, q), a) &&& \text{(A7-i), Skolemized} \\
\mathbf{B}(a, b, c) \rightarrow \mathbf{B}(c, b, a) &&& \text{(A14-i), symmetry of betweenness} \\
\mathbf{B}(a, b, d) \wedge \mathbf{B}(b, c, d) \rightarrow \mathbf{B}(a, b, c) &&& \text{(A15-i), inner transitivity} \\
\text{Col}(u, v, p) \wedge u \neq v \wedge \mathbf{T}(a, p, b) &\rightarrow && \text{two-point line-circle} \\
\exists y, z (az = ab \wedge ay = ab \wedge \mathbf{T}(y, p, z) \wedge (p \neq a \rightarrow y \neq z)) &&& 
\end{aligned}$$

The axioms of stability are as follows:

$$\begin{aligned}
\neg \neg \mathbf{B}(a, b, c) &\rightarrow \mathbf{B}(a, b, c) \\
\neg \neg E(a, b, c, d) &\rightarrow E(a, b, c, d) \\
\neg a \neq b &\rightarrow a = b
\end{aligned}$$

For reference we also state the circle-circle continuity principle, which is not an axiom but a theorem. The circles must have distinct centers but one of them could be a null circle (zero radius).

$$\begin{aligned}
ap = ax \wedge aq = az \wedge cp = cd \wedge cq = cd \wedge \\
\mathbf{T}(a, x, b) \wedge \mathbf{T}(a, b, z) \wedge a \neq b \rightarrow \\
\exists z_1, z_2 (bz_1 = bd \wedge az_1 = ab \wedge bz_2 = ab \wedge az_2 = ab)
\end{aligned}$$

**15.3. Ruler-and-compass Tarski geometry.** This theory uses **LPT** (logic of partial terms) as given in [2], p. 97. Its axioms are similar to intuitionistic Tarski geometry with Skolem functions, except that there is an additional 4-ary function symbol  $il$  with the axioms

$$\begin{aligned}
\text{Col}(a, b, x) \wedge \text{Col}(p, q, x) \wedge \neg (\text{Col}(a, b, p) \wedge \text{Col}(a, b, q)) &\rightarrow \\
x = il(a, b, p, q) &&& \text{Axiom } il\text{-i} \\
il(a, b, p, q) \downarrow \rightarrow \text{Col}(a, b, il(a, b, p, q)) \wedge \text{Col}(p, q, il(a, b, p, q)) &&& \text{Axiom } il\text{-ii}
\end{aligned}$$

The Skolem term  $\text{ip}(a, p, c, b, q)$  is replaced in the Skolemized inner Pasch axiom by  $il(a, q, b, p)$ . Point  $c$  does not occur in this term. The term  $\text{center}(a, b, c)$  in the triangle circumscription axiom is not changed.

#### REFERENCES

- [1] JEREMY AVIGAD, EDWARD DEAN, and JOHN MUMMA, *A formal system for Euclid's Elements*, *Review of Symbolic Logic*, vol. 2 (2009), pp. 700–768.
- [2] MICHAEL BEESON, *Foundations of constructive mathematics*, Springer-Verlag, Berlin Heidelberg New York, 1985.
- [3] ———, *Constructive geometry, Proceedings of the Tenth Asian Logic Colloquium, Kobe, Japan, 2008* (Singapore) (Tosiyasu Arai, editor), World Scientific, 2009, pp. 19–84.
- [4] ———, *Proof and computation in geometry, Automated deduction in geometry (ADG 2012)* (Tetsuo Ida and Jacques Fleuriot, editors), Springer Lecture Notes in Artificial Intelligence, vol. 7993, Springer, 2013, pp. 1–30.
- [5] ———, *Constructive geometry and the parallel postulate*, *The Bulletin of Symbolic Logic*, (submitted).

- [6] KAROL BORSUK and WANDA SZMIELEW, *Foundations of geometry: Euclidean and Bolyai-Lobachevskian geometry, projective geometry*, North-Holland, Amsterdam, 1960, translated from Polish by Erwin Marquit.
- [7] GABRIEL BRAUN and JULIEN NARBOUX, *From Tarski to Hilbert, Automated deduction in geometry 2012* (Tetsuo Ida and Jacques Fleuriot, editors), 2012.
- [8] L. E. J. BROUWER, *Contradictoriness of elementary geometry*, *L. E. J. Brouwer, Collected Works* (Arend Heyting, editor), North-Holland, 1975, pp. 497–498.
- [9] KURT GÖDEL, *Zur intuitionistischen arithmetik und zahlentheorie*, *Kurt Gödel, Collected Works volume I*, Oxford University Press, 1933, with English translation, pp. 286–295.
- [10] MARVIN JAY GREENBERG, *Euclidean and non-Euclidean geometries: Development and history*, fourth ed., W. H. Freeman, New York, 2008.
- [11] HARAGAURI NARAYAN GUPTA, *Contributions to the axiomatic foundations of geometry, Ph.D. thesis*, University of California, Berkeley, 1965.
- [12] ROBIN HARTSHORNE, *Geometry: Euclid and beyond*, Springer, 2000.
- [13] DAVID HILBERT, *Foundations of geometry (Grundlagen der Geometrie)*, Open Court, La Salle, Illinois, 1960, Second English edition, translated from the tenth German edition by Leo Unger. Original publication date, 1899.
- [14] STEPHEN C. KLEENE, Permutability of inferences in Gentzen’s calculi LK and LJ, *A.M.S. Memoirs*, vol. 10, pp. 1–26, *A.M.S. Memoirs*, American Mathematical Society, Providence, R. I., 1952, pp. 1–26.
- [15] T. J. M. MAKARIOS, *A further simplification of Tarski’s axioms of geometry*, *Note di Matematica*, vol. 33 (2013), no. 2, pp. 123–132.
- [16] MORITZ PASCH, *Vorlesung über Neuere Geometrie*, Teubner, Leipzig, 1882.
- [17] ZENON PIESYK, *Remarks on the axiomatic geometry of Tarski*, *Prace Mat.*, vol. 9 (1965), pp. 25–33.
- [18] J. F. RIGBY, *Congruence axioms for absolute geometry*, *Mathematical Chronicle*, vol. 4 (1975), pp. 13–44.
- [19] W. SCHWABHÄUSER, WANDA SZMIELEW, and ALFRED TARSKI, *Metamathematische Methoden in der Geometrie: Teil I: Ein axiomatischer Aufbau der euklidischen Geometrie. Teil II: Metamathematische Betrachtungen (Hochschultext)*, Springer-Verlag, 1983, Reprinted 2012 by Ishi Press, with a new foreword by Michael Beeson.
- [20] JACOB STEINER, *Einige geometrische Betrachtungen*, *Crelle’s Journal*, vol. I (1826), pp. 161–184 and 252–288, reprinted in [21], pp. 17–76.
- [21] ———, *Gesammelte Werke*, Chelsea Publishing Company, 1971.
- [22] J. STROMMER, *über die Kreisaxiome*, *Periodica Mathematica Hungarica*, vol. 4 (1973), pp. 3–16.
- [23] WANDA SZMIELEW, *Some metamathematical problems concerning elementary hyperbolic geometry*, *The axiomatic method with special reference to geometry and physics, proceedings of an international symposium, held at the University of California, Berkeley, December 26, 1957-January 4, 1958*. (Amsterdam) (Leon Henkin, Patrick Suppes, and Alfred Tarski, editors), *Studies in logic and the foundations of mathematics*, North-Holland, 1959, pp. 30–52.
- [24] ALFRED TARSKI, *What is elementary geometry?*, *The axiomatic method, with special reference to geometry and physics. proceedings of an international symposium held at the Univ. of Calif., Berkeley, Dec. 26, 1957-Jan. 4, 1958* (Amsterdam) (Leon Henkin, Patrick Suppes, and Alfred Tarski, editors), *Studies in Logic and the Foundations of Mathematics*, North-Holland, 1959, Available as a 2007 reprint, Brouwer Press, ISBN 1-443-72812-8, pp. 16–29.
- [25] ALFRED TARSKI and STEVEN GIVANT, *Tarski’s system of geometry*, *The Bulletin of Symbolic Logic*, vol. 5 (1999), no. 2, pp. 175–214.